

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/16, C07K 14/59, A61K 38/24	A1	(11) International Publication Number: WO 97/42322 (43) International Publication Date: 13 November 1997 (13.11.97)
(21) International Application Number: PCT/US96/06483 (22) International Filing Date: 8 May 1996 (08.05.96) (71) Applicant (for all designated States except US): THE GOVERNMENT OF THE UNITED STATES OF AMERICA as represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; National Institutes of Health, Office of Technology Transfer, Suite 325, 6011 Executive Boulevard, Rockville, MD 20852 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): SZKUDLINSKI, Mariusz, W. [PL/US]; 9738 Pleasant Gate Lane, Potomac, MD 20854 (US). WEINTRAUB, Bruce, D. [US/US]; 6620 Sulky Lane, Rockville, MD 20852 (US). GROSSMAN, Mathis [DE/US]; Apartment C902, 5901 Montrose Road, Rockville, MD 20852 (US). (74) Agents: NEEDLE, William, H. et al.; Needle & Rosenberg, Suite 1200, 127 Peachtree Street, N.E., Atlanta, GA 30303-1811 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>

(54) Title: GLYCOPROTEIN HORMONE SUPERAGONISTS**(57) Abstract**

The invention is directed toward a human glycoprotein hormone having at least one, two, three, four or five basic amino acids in the α -subunit at positions selected from the group consisting of positions 11, 13, 14, 16, 17 and 20. The invention is also directed to a human glycoprotein where at least one of the amino acids at positions 58, 63 and 69 of the β -subunit of the human thyroid stimulating hormone are basic amino acids. The invention is further directed to a modified human glycoprotein hormone having increased activity over a wild-type human glycoprotein hormone, where the modified human glycoprotein hormone comprises a basic amino acid substituted at a position corresponding to the same amino acid position in a non-human glycoprotein hormone having an increased activity over the wild-type human glycoprotein hormone. The invention is also directed to a method of constructing superactive nonchimeric analogs of human hormones comprising comparing the amino acid sequence of a more active homolog from another species to the human hormone, substituting selected amino acids in the human hormone with the corresponding amino acids from the other species, determining the activity of the substituted human hormones, and selecting superactive analogs from the substituted human hormones. The invention is also directed to nucleic acids encoding the modified human glycoprotein hormones, vectors containing those nucleic acids, and host cells containing those vectors.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

GLYCOPROTEIN HORMONE SUPERAGONISTS

BACKGROUND OF THE INVENTION

5 Field of the Invention

This invention relates generally to modified glycoprotein hormones. Specifically, this invention relates to modifications to a human glycoprotein which create superagonist activity.

10

Background Art

Thyrotropin (thyroid-stimulating hormone, TSH) and the gonadotropins chorionic gonadotropin, (CG), lutropin (luteinizing hormone, LH), and follitropin
15 (follicle-stimulating hormone, FSH) comprise the family of glycoprotein hormones. Each hormone is a heterodimer of two non-covalently linked subunits: α and β . Within the same species, the amino acid sequence of the α -subunit is identical in all the hormones, whereas the sequence of the β -subunit is hormone specific. (*Pierce, J.G. and Parsons, T.F.* "Glycoprotein hormones: structure and function." *Ann. Rev. Biochem.*
20 50:465-495 (1981)). The fact that the sequences of the subunits are highly conserved from fish to mammals implies that these hormones have evolved from a common ancestral protein (*Fontaine Y-A. and Burzawa-Gerard, E.* "Esquisse de l'evolution des hormones gonadotropes et thyreotropes des vertebres." *Gen. Comp. Endocrinol.* 32:341-347 (1977)). Evolutionary changes of these hormones resulted in certain cases in
25 modification of biological activity (*Licht, P. et al.* "Evolution of gonadotropin structure and function." *Rec. Progr. Horm. Res.*, 33:169-248 (1977) and *Combarnous, Y.* "Molecular basis of the specificity of binding of glycoprotein hormones to their receptors." *Endocrine Rev.* 13:670-691 (1992)), although, specific structural determinants modulating biopotency have not been elucidated. For example, human
30 thyroid stimulating hormone (hTSH) and bovine thyroid stimulating hormone (bTSH) share high homology in the α (70%) and β (89%) subunit sequence, but bTSH is 6-10 fold more potent than hTSH (*Yamazaki, K. et al.* "Potent thyrotropic activity of human chorionic gonadotropin variants in terms of ^{125}I incorporation and de novo synthesized

thyroid hormone release in human thyroid follicles." J. Clin. Endocrinol. Metab. 80:473-479 (1995)).

Glycoprotein hormones are crucial in certain therapies, such as in the treatment
5 of patients with thyroid carcinoma. (See, for example, *Meier, C.A., et al.*, "Diagnostic
use of Recombinant Human Thyrotropin in Patients with Thyroid Carcinoma (Phase I/II
Study)." J. Clin. Endocrinol. Metabol. 78:22 (1994)). The potential use of human
thyroid stimulating hormone (TSH) in the treatment of this disease has been abandoned
due to the potential transmission of Creutzfeldt-Jakob disease. An alternative to the use
10 of human TSH is the use of bovine TSH, but this approach is very limited since this
hormone causes side-effects such as nausea, vomiting, local induration, urticaria, and a
relatively high possibility of anaphylactic shock (*Meier, C.A., et al.*). The lack of
bioconsistency of urinary gonadotropins and the limited efficacy of recombinant
glycoprotein hormones justify their further replacement with more effective recombinant
15 analogs. Therefore, there is a need for human-derived glycoprotein hormones as well as
agonists of these hormones.

For example the administration of an agonist of the thyroid stimulating hormone
in a particular clinical situation such as thyroid carcinoma, will enhance the uptake of
20 radioiodine into the carcinoma to treat the disease. Agonists of the thyroid stimulating
hormone will cause a greater amount of the radioiodine to be targeted to the carcinoma,
thereby resulting in a more effective treatment. Alternatively, glycoprotein hormones
used to induce ovulation can be replaced with superagonists. This will lower the
required dose of the hormone which currently is a major medical problem in fertility
25 treatment. (*Ben-Rafael, Z., et al.* "Pharmacokinetics of follicle-stimulating hormone:
clinical significance." Fertility and Sterility 63:689 (1995)). Where the use of wild-type
follicle stimulating hormone has led to hyperstimulation and higher rates of multiple
pregnancies and abortions, apparently by a high number of hormone molecules
stimulating many follicles, a superagonist of follicle-stimulating hormone can be
30 administered to treat the infertility. The use of an agonist of this modified hormone can

result in a lower frequency of stimulation of multiple follicles since a lower number of hormone molecules can be administered to achieve the desired result.

The present invention provides, for the first time, specific amino acid
5 substitutions in human glycoprotein hormones which results in human glycoprotein hormone analogs that show a major increase in both *in vitro* and *in vivo* bioactivity.

SUMMARY OF THE INVENTION

10 In accordance with the purpose(s) of this invention, as embodied and broadly described herein, this invention, in one aspect, provides a human glycoprotein hormone comprising at least three basic amino acids in the α -subunit at positions selected from the group consisting of positions 11, 13, 14, 16, 17 and 20.

15 The invention further provides a human glycoprotein hormone comprising at least one basic amino acid in the α -subunit at positions selected from the group consisting of positions 11, 13, 14, 16, 17 and 20.

20 In another aspect, the invention provides a modified human glycoprotein hormone having increased activity over a wild-type human glycoprotein, wherein the modified human hormone comprises a basic amino acid substituted at a position corresponding to the same amino acid position in a non-human glycoprotein hormone having an increased activity over the wild-type human glycoprotein.

25 In another aspect, the invention provides a method of treating a condition associated with a glycoprotein hormone activity in a subject comprising administering a therapeutic amount of the glycoprotein hormone of the present invention to the patient.

30 In another aspect, the invention provides a method of constructing superactive nonchimeric analogs of human hormones comprising comparing the amino acid sequence of a more active homolog from another species to the human hormone,

substituting amino acids in the human hormone with the corresponding amino acids from the other species, determining the activity of the substituted human hormone, and selecting superactive analogs from the substituted human hormones.

- 5 In yet another aspect, the present invention provides nucleic acids which encode the modified glycoprotein hormones.

BRIEF DESCRIPTION OF THE DRAWINGS

- 10 Fig. 1 shows a comparison of the relevant primary sequences of the α -subunit from 27 different species (a). Alignment of the subunit sequences obtained from sequencing of PCR amplified fragment of genomic DNA in chimpanzee, orangutan, gibbon and baboon (underlined), received from GeneBank, SWISS-PROT and PDB databank were made. The numbering of the sequences corresponds to that of human α -
- 15 subunit sequence. Dashes (---) indicate amino acid residues which are identical to those of the human α -subunit. Conserved among different species lysine residues are bolded. The primate sequences determined in this study are underlined. The human, chimpanzee and orangutan α -subunit sequences are the only sequences without basic amino acids in this region, despite the relatively high degree of similarity in diverse vertebrate species.
- 20 (b) Mutations of human sequence made in this region included introduction of single and multiple Lys residues present in all non-human mammalian sequences. Additionally, alanine mutagenesis of residues 13, 16 and 20 was used to study the role of Gln13, Pro16 and Gln20.
- 25 Fig. 2 shows the bioactivities and receptor binding activities of the most potent hTSH analogs: (a, b) cAMP stimulation in CHO-JP09 cells. Data represent the mean \pm SEM of triplicate determinations from a representative experiment repeated three (a) and two (b) times. (c, d) Receptor-binding activities to CHO-JP09 cells. The same mutants tested as in the Fig. 2a and Fig. 2b respectively. Values are the mean \pm SEM of
- 30 quadruplicate determinations from one experiment, repeated two times. (e) Thymidine uptake stimulation in FRTL-5 cells. Values are the mean \pm SEM of quadruplicate

determinations from one experiment, repeated two times. (f) Stimulation of T_4 secretion in mice. Each data point represents the mean \pm SEM of values from 4-5 animals of a representative experiment repeated two times. (g) cAMP stimulation in CHO-hTSH cells. Data represent the mean \pm SEM of 3-4 determinations from a representative
5 experiment repeated 3 times. (h) Receptor-binding activities in CHO-JP09 cells. Data represent the mean \pm SEM of 3-4 determinations from a representative experiment repeated 3 times. (i) Stimulation of T_4 secretion in mice. Each data point represents the mean \pm SEM of values from 4-5 animals of a representative experiment repeated two times.

10

Fig. 3 shows the bioactivities and receptor binding activities of the most potent hCG analogs. Progesterone production stimulation (a) and receptor binding assay (b) in MA-10 cells. Data represent the mean \pm SEM of triplicate determinations from a representative experiment repeated three times. The relative maximal production levels
15 of progesterone are presented in the Table II as % obtained with WT-hCG. cAMP stimulation (c) and receptor binding assay (d) in COS-7 cells expressing hLH receptor. Data represent the mean \pm SEM of triplicate determinations from a representative experiment repeated two times.

20

DETAILED DESCRIPTION OF THE INVENTION

The present invention may be understood more readily by reference to the following detailed description of the preferred embodiments of the invention and the
25 Example included therein and to the Figures and their previous and following description.

Before the present compounds, compositions, and methods are disclosed and described, it is to be understood that this invention is not limited to specific hormones,
30 specific subjects, i.e. humans as well as non-human mammals, specific amino acids, specific clinical conditions, specific analogs, or specific methods, as such may, of course,

vary, and the numerous modifications and variations therein will be apparent to those skilled in the art. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

5 As used in the specification and in the claims, "a" can mean one or more, depending upon the context in which it is used. Thus, for example, reference to "a human glycoprotein hormone" means that at least one human glycoprotein hormone is utilized.

10 In one aspect, the invention provides a human glycoprotein hormone comprising at least three basic amino acids in the α -subunit at positions selected from the group consisting of positions 11, 13, 14, 16, 17 and 20.

 The invention further provides a human glycoprotein hormone comprising at
15 least one basic amino acid in the α -subunit at positions selected from the group consisting of positions 11, 13, 14, 16, 17 and 20.

 In another aspect, the invention provides a modified human glycoprotein hormone having increased activity over a wild-type human glycoprotein, wherein the
20 modified human hormone comprises a basic amino acid substituted at a position corresponding to the same amino acid position in a non-human glycoprotein hormone having an increased activity over the wild-type human glycoprotein.

 In another aspect, the invention provides a method of treating a condition
25 associated with a glycoprotein hormone activity in a subject comprising administering a therapeutic amount of the glycoprotein hormone of the present invention to the patient.

 In another aspect, the invention provides a method of assisting reproduction in a subject comprising administering an assisting amount of the glycoprotein hormone of the
30 present invention.

In another aspect, the invention provides a method of constructing superactive nonchimeric analogs of human hormones comprising comparing the amino acid sequence of a more active homolog from another species to the human hormone, substituting amino acids in the human hormone with the corresponding amino acids from the other species, determining the activity of the substituted human hormone, and selecting superactive analogs from the substituted human hormones.

By "human" glycoprotein hormone is meant that the number of amino acid substitutions made in the wild-type sequence does not exceed one-half the number of amino acid differences at corresponding positions in the corresponding polypeptide hormones between human and another species. Thus, the modified polypeptide hormone would be considered more like the wild-type polypeptide hormone of the human than the corresponding polypeptide hormone from the non-human species from which the amino acid substitutions are derived, based on the amino acid coding sequence. For example, if there were a total of 20 amino acid differences at corresponding positions in corresponding glycoprotein hormones between a human glycoprotein and a bovine glycoprotein hormone, a "human" glycoprotein hormone would be a modified wild-type human hormone which contains 10 or fewer amino acid substitutions within its amino acid sequence which are homologous to the corresponding amino acids in the bovine amino acid sequence. More specifically, the thyroid stimulating hormone, as set forth in the Examples contained herein, would be considered "human" if 20 or more of the 40 total amino acid differences between the α - and β -subunits of the human and the bovine homologs are homologous to the amino acid at the corresponding position in the human thyroid stimulating hormone.

25

Naturally, because of the risk of an adverse immune response to the administration of the modified glycoprotein hormone where the recipient of the modified glycoprotein hormone is a human, the modified glycoprotein hormone is preferably homologous to the human amino acid sequence to the greatest extent possible without an unacceptable loss in the superagonist activity. Alternatively, where the subject being administered the modified glycoprotein is non-human, the modified glycoprotein

30

hormone is preferably homologous to the specific non-human amino acid sequence to the greatest extent possible without an unacceptable loss in the superagonist activity. Thus, in a preferred embodiment of the present invention, in modifying a wild-type glycoprotein to construct a modified glycoprotein with a superagonist activity by
5 substituting specific amino acids, the substituted amino acids which do not increase agonist activity number 10 or less, especially 9, 8, 7, 6, 5, 4, 3, and 2 or zero.

Likewise, by "nonchimeric" is meant that the number of amino substitutions does not exceed one-half the number of amino acid differences at corresponding positions in
10 the corresponding polypeptide hormones between species, such that the modified polypeptide hormone would be considered more like the wild-type polypeptide hormone of the species being modified than the corresponding polypeptide hormone from the species from which the amino acid substitutions are derived, based on the amino acid coding sequence.

15

In yet another aspect, the present invention provides nucleic acids which encode the modified glycoprotein hormones.

Glycoprotein hormones comprise a family of hormones which are structurally
20 related heterodimers consisting of a species-common α -subunit and a distinct β -subunit that confers the biological specificity for each hormone. For a general review of glycoprotein hormones, see *Pierce, J.G. et al.*, " Glycoprotein hormones: structure and function." *Ann. Rev. Biochem.* 50:465-495 (1981), see also *Combarnous, Y.* "Molecular basis of the specificity of binding of glycoprotein hormones to their receptors."
25 *Endocrine Rev.* 13:670-691 (1992) This family of hormones includes chorionic gonadotropin (CG), lutropin (luteinizing hormone, LH), follitropin (follicle-stimulating hormone, FSH), and thyrotropin (thyroid-stimulating hormone, TSH). Each of these glycoprotein hormones with at least one basic amino acid in the α -subunit at positions selected from the group consisting of positions 11, 13, 14, 16, 17, and 20, is provided by
30 the present invention.

Basic amino acids comprise the amino acids lysine, arginine, and histidine, and any other basic amino acid which may be a modification to any of these three amino acids, synthetic basic amino acids not normally found in nature, or any other amino acid which is positively charged at a neutral pH.

5

The glycoprotein hormones provided for by the present invention may be obtained in any number of ways. For example, a DNA molecule encoding a glycoprotein hormone can be isolated from the organism in which it is normally found. For example, a genomic DNA or cDNA library can be constructed and screened for the
10 presence of the nucleic acid of interest. Methods of constructing and screening such libraries are well known in the art and kits for performing the construction and screening steps are commercially available (for example, Stratagene Cloning Systems, La Jolla, CA). Once isolated, the nucleic acid can be directly cloned into an appropriate vector, or if necessary, be modified to facilitate the subsequent cloning steps. Such modification
15 steps are routine, an example of which is the addition of oligonucleotide linkers which contain restriction sites to the termini of the nucleic acid. General methods are set forth in *Sambrook et al.*, "Molecular Cloning, a Laboratory Manual," Cold Spring Harbor Laboratory Press (1989).

20 Once the nucleic acid sequence of the desired glycoprotein hormone is obtained, basic amino acids can be positioned at any particular amino acid positions by techniques well known in the art. For example, PCR primers can be designed which span the amino acid position or positions and which can substitute a basic amino acid for a non-basic amino acid. Then a nucleic acid can be amplified and inserted into the wild-type
25 glycoprotein hormone coding sequence in order to obtain any of a number of possible combinations of basic amino acids at any position of the glycoprotein hormone. Alternatively, one skilled in the art can introduce specific mutations at any point in a particular nucleic acid sequence through techniques for point mutagenesis. General methods are set forth in *Smith, M.* "In vitro mutagenesis" *Ann. Rev. Gen.*, 19:423-462
30 (1985) and *Zoller, M.J.* "New molecular biology methods for protein engineering" *Curr. Opin. Struct. Biol.*, 1:605-610 (1991).

Another example of a method of obtaining a DNA molecule encoding a specific glycoprotein hormone is to synthesize a recombinant DNA molecule which encodes the glycoprotein hormone. For example, oligonucleotide synthesis procedures are routine in the art and oligonucleotides coding for a particular protein region are readily obtainable through automated DNA synthesis. A nucleic acid for one strand of a double-stranded molecule can be synthesized and hybridized to its complementary strand. One can design these oligonucleotides such that the resulting double-stranded molecule has either internal restriction sites or appropriate 5' or 3' overhangs at the termini for cloning into an appropriate vector. Double-stranded molecules coding for relatively large proteins can readily be synthesized by first constructing several different double-stranded molecules that code for particular regions of the protein, followed by ligating these DNA molecules together. For example, *Cunningham, et al.*, "Receptor and Antibody Epitopes in Human Growth Hormone Identified by Homolog-Scanning Mutagenesis," *Science*, 243:1330-1336 (1989), have constructed a synthetic gene encoding the human growth hormone gene by first constructing overlapping and complementary synthetic oligonucleotides and ligating these fragments together. See also, *Ferretti, et al.*, *Proc. Nat. Acad. Sci.* 82:599-603 (1986), wherein synthesis of a 1057 base pair synthetic bovine rhodopsin gene from synthetic oligonucleotides is disclosed. By constructing a glycoprotein hormone in this manner, one skilled in the art can readily obtain any particular glycoprotein hormone with basic amino acids at any particular position or positions of either the α -subunit, the β -subunit, or both. See also, U.S. Patent No. 5,503,995 which describes an enzyme template reaction method of making synthetic genes. Techniques such as this are routine in the art and are well documented. DNA fragments encoding glycoprotein hormones can then be expressed *in vivo* or *in vitro* as discussed below.

Once a nucleic acid encoding a particular glycoprotein hormone of interest, or a region of that nucleic acid, is constructed, modified, or isolated, that nucleic acid can then be cloned into an appropriate vector, which can direct the *in vivo* or *in vitro* synthesis of that wild-type and/or modified glycoprotein hormone. The vector is

contemplated to have the necessary functional elements that direct and regulate transcription of the inserted gene, or hybrid gene. These functional elements include, but are not limited to, a promoter, regions upstream or downstream of the promoter, such as enhancers that may regulate the transcriptional activity of the promoter, an
5 origin of replication, appropriate restriction sites to facilitate cloning of inserts adjacent to the promoter, antibiotic resistance genes or other markers which can serve to select for cells containing the vector or the vector containing the insert, RNA splice junctions, a transcription termination region, or any other region which may serve to facilitate the expression of the inserted gene or hybrid gene. (See generally, Sambrook *et al.*).

10

There are numerous *E. coli* (*Escherichia coli*) expression vectors known to one of ordinary skill in the art which are useful for the expression of the nucleic acid insert. Other microbial hosts suitable for use include bacilli, such as *Bacillus subtilis*, and other enterobacteriaceae, such as *Salmonella*, *Serratia*, and various *Pseudomonas* species. In
15 these prokaryotic hosts one can also make expression vectors, which will typically contain expression control sequences compatible with the host cell (e.g., an origin of replication). In addition, any number of a variety of well-known promoters will be present, such as the lactose promoter system, a tryptophan (Trp) promoter system, a beta-lactamase promoter system, or a promoter system from phage lambda. The
20 promoters will typically control expression, optionally with an operator sequence, and have ribosome binding site sequences for example, for initiating and completing transcription and translation. If necessary, an amino terminal methionine can be provided by insertion of a Met codon 5' and in-frame with the downstream nucleic acid insert. Also, the carboxy-terminal extension of the nucleic acid insert can be removed
25 using standard oligonucleotide mutagenesis procedures.

Additionally, yeast expression can be used. There are several advantages to yeast expression systems. First, evidence exists that proteins produced in a yeast secretion systems exhibit correct disulfide pairing. Second, post-translational
30 glycosylation is efficiently carried out by yeast secretory systems. The *Saccharomyces cerevisiae* pre-pro-alpha-factor leader region (encoded by the *MF"-I* gene) is routinely

used to direct protein secretion from yeast. (*Brake, et al.*, " α -Factor-Directed Synthesis and Secretion of Mature Foreign Proteins in *Saccharomyces cerevisiae*." *Proc. Nat. Acad. Sci.*, 81:4642-4646 (1984)). The leader region of pre-pro-alpha-factor contains a signal peptide and a pro-segment which includes a recognition sequence for a yeast

5 protease encoded by the *KEX2* gene: this enzyme cleaves the precursor protein on the carboxyl side of a Lys-Arg dipeptide cleavage signal sequence. The nucleic acid coding sequence can be fused in-frame to the pre-pro-alpha-factor leader region. This construct is then put under the control of a strong transcription promoter, such as the alcohol dehydrogenase I promoter or a glycolytic promoter. The nucleic acid coding sequence

10 is followed by a translation termination codon which is followed by transcription termination signals. Alternatively, the nucleic acid coding sequences can be fused to a second protein coding sequence, such as Sj26 or β -galactosidase, used to facilitate purification of the fusion protein by affinity chromatography. The insertion of protease cleavage sites to separate the components of the fusion protein is applicable to

15 constructs used for expression in yeast. Efficient post translational glycosylation and expression of recombinant proteins can also be achieved in Baculovirus systems.

Mammalian cells permit the expression of proteins in an environment that favors important post-translational modifications such as folding and cysteine pairing, addition

20 of complex carbohydrate structures, and secretion of active protein. Vectors useful for the expression of active proteins in mammalian cells are characterized by insertion of the protein coding sequence between a strong viral promoter and a polyadenylation signal. The vectors can contain genes conferring hygromycin resistance, gentamicin resistance, or other genes or phenotypes suitable for use as selectable markers, or methotrexate

25 resistance for gene amplification. The chimeric protein coding sequence can be introduced into a Chinese hamster ovary (CHO) cell line using a methotrexate resistance-encoding vector, or other cell lines using suitable selection markers. Presence of the vector DNA in transformed cells can be confirmed by Southern blot analysis. Production of RNA corresponding to the insert coding sequence can be confirmed by

30 Northern blot analysis. A number of other suitable host cell lines capable of secreting intact human proteins have been developed in the art, and include the CHO cell lines,

HeLa cells, myeloma cell lines, Jurkat cells, etc. Expression vectors for these cells can include expression control sequences, such as an origin of replication, a promoter, an enhancer, and necessary information processing sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences.

- 5 Preferred expression control sequences are promoters derived from immunoglobulin genes, SV40, Adenovirus, Bovine Papilloma Virus, etc. The vectors containing the nucleic acid segments of interest can be transferred into the host cell by well-known methods, which vary depending on the type of cellular host. For example, calcium chloride transformation is commonly utilized for prokaryotic cells, whereas calcium
10 phosphate, DEAE dextran, or lipofectin mediated transfection or electroporation may be used for other cellular hosts.

- Alternative vectors for the expression of genes in mammalian cells, those similar to those developed for the expression of human gamma-interferon, tissue plasminogen
15 activator, clotting Factor VIII, hepatitis B virus surface antigen, protease NexinI, and eosinophil major basic protein, can be employed. Further, the vector can include CMV promoter sequences and a polyadenylation signal available for expression of inserted nucleic acids in mammalian cells (such as COS-7).

- 20 Expression of the gene or hybrid gene can be by either *in vivo* or *in vitro*. *In vivo* synthesis comprises transforming prokaryotic or eukaryotic cells that can serve as host cells for the vector. An example of modified glycoprotein hormones inserted into a prokaryotic expression vector is given in the Example section contained herein.

- 25 Alternatively, expression of the gene can occur in an *in vitro* expression system. For example, *in vitro* transcription systems are commercially available which are routinely used to synthesize relatively large amounts of mRNA. In such *in vitro* transcription systems, the nucleic acid encoding the glycoprotein hormone would be cloned into an expression vector adjacent to a transcription promoter. For example, the
30 Bluescript II cloning and expression vectors contain multiple cloning sites which are flanked by strong prokaryotic transcription promoters. (Stratagene Cloning Systems, La

Jolla, CA). Kits are available which contain all the necessary reagents for *in vitro* synthesis of an RNA from a DNA template such as the Bluescript vectors. (Stratagene Cloning Systems, La Jolla, CA). RNA produced *in vitro* by a system such as this can then be translated *in vitro* to produce the desired glycoprotein hormone. (Stratagene
5 Cloning Systems, La Jolla, CA).

Another method of producing a glycoprotein hormone is to link two peptides or polypeptides together by protein chemistry techniques. For example, peptides or polypeptides can be chemically synthesized using currently available laboratory
10 equipment using either Fmoc (9-fluorenylmethyloxycarbonyl) or Boc (*tert*-butyloxycarbonyl) chemistry. (Applied Biosystems, Inc., Foster City, CA). One skilled in the art can readily appreciate that a peptide or polypeptide corresponding to a hybrid glycoprotein hormone can be synthesized by standard chemical reactions. For example, a peptide or polypeptide can be synthesized and not cleaved from its synthesis
15 resin whereas the other fragment of a hybrid peptide can be synthesized and subsequently cleaved from the resin, thereby exposing a terminal group which is functionally blocked on the other fragment. By peptide condensation reactions, these two fragments can be covalently joined via a peptide bond at their carboxyl and amino termini, respectively, to form a hybrid peptide. (*Grant, G.A., "Synthetic Peptides: A*
20 *User Guide," W.H. Freeman and Co., N.Y. (1992) and Bodansky, M. and Trost, B., Ed., "Principles of Peptide Synthesis," Springer-Verlag Inc., N.Y. (1993)*). Alternatively, the peptide or polypeptide can be independently synthesized *in vivo* as described above. Once isolated, these independent peptides or polypeptides may be linked to form a glycoprotein hormone via similar peptide condensation reactions.

25

For example, enzymatic ligation of cloned or synthetic peptide segments can allow relatively short peptide fragments to be joined to produce larger peptide fragments, polypeptides or whole protein domains (*Abrahmsen, L., et al., Biochemistry, 30:4151 (1991)*). Alternatively, native chemical ligation of synthetic peptides can be
30 utilized to synthetically construct large peptides or polypeptides from shorter peptide fragments. This method consists of a two step chemical reaction (*Dawson, et al.,*

"Synthesis of Proteins by Native Chemical Ligation," *Science*, 266:776-779 (1994)).
The first step is the chemoselective reaction of an unprotected synthetic peptide- α -thioester with another unprotected peptide segment containing an amino-terminal Cys residue to give a thioester-linked intermediate as the initial covalent
5 product. Without a change in the reaction conditions, this intermediate undergoes spontaneous, rapid intramolecular reaction to form a native peptide bond at the ligation site. Application of this native chemical ligation method to the total synthesis of a protein molecule is illustrated by the preparation of human interleukin 8 (IL-8) (*Clark-Lewis, I., et al.*, FEBS Lett., 307:97 (1987), *Clark-Lewis, I., et al.*, J.Biol.Chem.,
10 269:16075 (1994), *Clark-Lewis, I., et al.*, Biochemistry, 30:3128 (1991), and *Rajaratnam, K., et al.*, Biochemistry, 29:1689 (1994)).

Alternatively, unprotected peptide segments can be chemically linked where the bond formed between the peptide segments as a result of the chemical ligation is an
15 unnatural (non-peptide) bond (*Schnolzer, M., et al.*, *Science*, 256:221 (1992)). This technique has been used to synthesize analogs of protein domains as well as large amounts of relatively pure proteins with full biological activity (*deLisle Milton, R.C., et al.*, "Techniques in Protein Chemistry IV," Academic Press, New York, pp. 257-267 (1992)).

20

The invention also provides fragments of modified glycoprotein hormones which have either superagonist or antagonist activity. The polypeptide fragments of the present invention can be recombinant proteins obtained by cloning nucleic acids encoding the polypeptide in an expression system capable of producing the polypeptide
25 fragments thereof. For example, one can determine the active domain of a glycoprotein hormone which, together with a β -subunit, can interact with a glycoprotein hormone receptor and cause a biological effect associated with the glycoprotein hormone. In one example, amino acids found to not contribute to either the activity or the binding specificity or affinity of the glycoprotein hormone can be deleted without a loss in the
30 respective activity.

For example, amino or carboxy-terminal amino acids can be sequentially removed from either the native or the modified glycoprotein hormone and the respective activity tested in one of many available assays. In another example, a fragment of a modified glycoprotein can comprise a modified hormone wherein at least one amino acid has been substituted for the naturally occurring amino acid at specific positions in either the α or the β -subunit, and a portion of either amino terminal or carboxy terminal amino acids, or even an internal region of the hormone, has been replaced with a polypeptide fragment or other moiety, such as biotin, which can facilitate in the purification of the modified glycoprotein hormone. For example, a modified glycoprotein can be fused to a maltose binding protein, through either peptide chemistry or cloning the respective nucleic acids encoding the two polypeptide fragments into an expression vector such that the expression of the coding region results in a hybrid polypeptide. The hybrid polypeptide can be affinity purified by passing it over an amylose affinity column, and the modified glycoprotein can then be separated from the maltose binding region by cleaving the hybrid polypeptide with the specific protease factor Xa. (See, for example, New England Biolabs Product Catalog, 1996, pg. 164.)

Active fragments of a glycoprotein hormone can also be synthesized directly or obtained by chemical or mechanical disruption of larger glycoprotein hormone. An active fragment is defined as an amino acid sequence of at least about 5 consecutive amino acids derived from the naturally occurring amino acid sequence, which has the relevant activity, e.g., binding or regulatory activity.

The fragments, whether attached to other sequences or not, can also include insertions, deletions, substitutions, or other selected modifications of particular regions or specific amino acids residues, provided the activity of the peptide is not significantly altered or impaired compared to the modified glycoprotein hormone. These modifications can provide for some additional property, such as to remove/add amino acids capable of disulfide bonding, to increase its bio-longevity, etc. In any case, the peptide must possess a bioactive property, such as binding activity, regulation of binding at the binding domain, etc. Functional or active regions of the glycoprotein

hormone may be identified by mutagenesis of a specific region of the hormone, followed by expression and testing of the expressed polypeptide. Such methods are readily apparent to a skilled practitioner in the art and can include site-specific mutagenesis of the nucleic acid encoding the receptor. (*Zoller, M.J. et al.*).

5

In one embodiment of the present invention, the human glycoprotein hormone comprises at least one basic amino acid in the α -subunit at the position selected from the group consisting of positions 11, 13, 14, 16, 17, and 20. In one embodiment, the human glycoprotein hormone has a basic amino acid at position 11. In another embodiment, 10 the human glycoprotein hormone has a basic amino acid at position 13. In another embodiment, the human glycoprotein hormone has a basic amino acid at position 14. In another embodiment, the human glycoprotein hormone has a basic amino acid at position 16. In another embodiment, the human glycoprotein hormone has a basic amino acid at position 17. In another embodiment, the human glycoprotein hormone has 15 a basic amino acid at position 20. In another embodiment of the present invention, the basic amino acid at position 11, 13, 14, 16, and 20 is lysine. In yet another embodiment of the present invention, the basic amino acid at position 17 is arginine.

The present invention also provides for a human glycoprotein hormone with 20 basic amino acids in the α -subunit in all combinations of any two positions selected from the group consisting of positions 11, 13, 14, 16, 17, and 20. For example, basic amino acids may be present at positions 11 and 13, or positions 11, and 14, or positions 11 and 16, or positions 11 and 17, or positions 11 and 20, or positions 13 and 14, or positions 13 and 17, or positions 14 and 16, or positions 14 and 17, or positions 14 and 20, or 25 positions 16 and 17, or positions 17 and 20. In one embodiment of the present invention, the human glycoprotein hormone has basic amino acids at position 16 and 13. In another embodiment of the present invention, the human glycoprotein hormone has basic amino acids at positions 20 and 13. In yet another embodiment, the human glycoprotein hormone has basic amino acids at positions 16 and 20.

30

The present invention also provides for a human glycoprotein hormone with basic amino acids in the α -subunit in all combinations of any three positions selected from the group consisting of positions 11, 13, 14, 16, 17, and 20. For example, basic amino acids may be present at positions 11, 13, and 14, or positions 11, 13, and 16, or positions 11, 13, and 17, or positions 11, 13, and 20, or positions 11, 14, and 16, or positions 11, 14, and 17, or positions 11, 14, and 20, or positions 11, 16, and 17, or positions 11, 16, and 20, or positions 11, 17, and 20, or positions 13, 14, and 16, or positions 13, 14, and 17, or positions 13, 14, and 20, or positions 13, 16, and 17, or positions 13, 17, and 20, or positions 14, 16, and 17, or positions 14, 16, and 20, or positions 14, 17, and 20, or positions 16, 17, and 20. In a preferred embodiment of the present invention, the human glycoprotein hormone has basic amino acids at positions 13, 16, and 20. In another embodiment of the present invention, the hormone is thyroid stimulating hormone. In another embodiment of the present invention, the hormone is follicle-stimulating hormone. In another embodiment of the present invention, the hormone is luteinizing hormone. In another embodiment of the present invention, the hormone is chorionic gonadotropin. In yet another embodiment of the present invention, the basic amino acids at any three positions selected from the group consisting of positions 11, 13, 14, 16, 17, and 20, are lysine.

20

The present invention also provides for a human thyroid stimulating hormone with at least three basic amino acids in the α -subunit at positions selected from the group consisting of positions 11, 13, 14, 16, 17, and 20, where the thyroid stimulating hormone also has a basic amino acid in at least one position selected from the group consisting of positions 58, 63, and 69 of the β -subunit. In one embodiment of the present invention, the thyroid stimulating hormone has a basic amino acid in at position 58 of the β -subunit. In another embodiment of the present invention, the thyroid stimulating hormone has a basic amino acid in at position 63 of the β -subunit. In another embodiment of the present invention, the thyroid stimulating hormone has a basic amino acid in at position 69 of the β -subunit. In another embodiment of the present invention, the thyroid stimulating hormone has a basic amino acid in each of

positions 58, 63, and 69 of the β -subunit. In yet another embodiment of the present invention, the basic amino acid in at least one position selected from the group consisting of positions 58, 63, and 69 of the β -subunit is arginine.

5 The present invention also provides for a human glycoprotein hormone with basic amino acids in the α -subunit in all combinations of any four positions selected from the group consisting of positions 11, 13, 14, 16, 17, and 20. One skilled in the art will readily determine the possible combinations available. In one embodiment, the human glycoprotein hormone has basic amino acids at positions 11, 13, 16, and 20. In
10 another embodiment, the human glycoprotein hormone has basic amino acids at positions 11, 13, 17, and 20. In another embodiment, the human glycoprotein hormone has basic amino acids at positions 13, 14, 17, and 20. In a preferred embodiment, the human glycoprotein hormone has basic amino acids at positions 13, 14, 16, and 20. In yet another embodiment of the present invention, the basic amino acids at any four
15 positions selected from the group consisting of positions 11, 13, 14, 16, 17, and 20, are lysine.

 The present invention also provides for a human glycoprotein hormone with basic amino acids in the α -subunit in all combinations of any five positions selected from
20 the group consisting of positions 11, 13, 14, 16, 17, and 20. One skilled in the art will readily determine the possible combinations available. In one embodiment, the human glycoprotein hormone has basic amino acids at positions 13, 14, 16, 17, and 20. In another embodiment, the human glycoprotein hormone has basic amino acids at positions 11, 13, 14, 16, and 20. In yet another embodiment of the present invention,
25 the basic amino acids at any five positions selected from the group consisting of positions 11, 13, 14, 16, 17, and 20, are selected from the group consisting of lysine and arginine.

 The present invention also provides for a human glycoprotein hormone with
30 basic amino acids in the α -subunit in all six of positions 11, 13, 14, 16, 17, and 20.

In another aspect, the present invention provides a human glycoprotein hormone with a basic amino acid in the α -subunit in at least one position selected from the group consisting of positions 11, 13, 14, 16, 17, and 20, wherein the hormone is human thyroid stimulating hormone and there is a basic amino acid in at least one position selected

5 from the group consisting of positions 58, 63, and 69 of the β -subunit. In one embodiment of the present invention, the human glycoprotein hormone has a basic amino acid at position 58 of the β -subunit of the human thyroid stimulating hormone. In another embodiment of the present invention, the human glycoprotein hormone has a basic amino acid at position 63 of the β -subunit of the human thyroid stimulating

10 hormone. In a preferred embodiment of the present invention, the human glycoprotein hormone has a basic amino acid at position 69 of the β -subunit of the human thyroid stimulating hormone. In another embodiment of the present invention, the human glycoprotein hormone has basic amino acids at position 58, 63, and 69 of the β -subunit of the human thyroid stimulating hormone. In yet another embodiment of the present

15 invention, the basic amino acid at the position selected from the group consisting of positions 58, 63, and 69 is arginine.

In another aspect, the present invention provides a human follicle-stimulating hormone, a human luteinizing hormone, or a human chorionic gonadotropin

20 glycoprotein hormone, wherein the hormone comprises a basic amino acid in at least one position selected from the group consisting of positions in the β -subunit of any of the glycoprotein hormones, corresponding to positions 58, 63, and 69 of the β -subunit of the human thyroid stimulating hormone. This approach applies equally to non-humans as well. For example, the β -subunit amino acid sequences of two bovine glycoprotein

25 hormones can be compared and substitutions made to any of the subunits based on the sequence differences.

One skilled in the art can readily determine which sites of the β -subunits of the other glycoprotein hormones correspond to sites 58, 63, and 69 of the β -subunit of the

30 human thyroid stimulating hormone. For example, see *Ward, et al.*, In: Bellet, D and Bidard, J.M. (eds) "Structure-function relationships of gonadotropins" Serono

Symposium Publications, Raven Press, New York, 65:1-19 (1990), where the amino acid sequences of 26 various glycoprotein hormone β -subunits are aligned and compared. Therefore, one skilled in the art can readily substitute non-basic amino acids at these sites of the other glycoprotein hormones for basic amino acids.

5

Similarly, the present invention provides for any human glycoprotein, wherein the hormone comprises a basic amino acid in at least one position selected from the group consisting of positions in the β -subunit of a glycoprotein hormone corresponding to the same positions in any of the other human glycoprotein hormones. For example, the amino acid sequence of the β -subunits of the human luteinizing hormone and the human chorionic gonadotropin hormone can be compared and amino acid substitutions made at selected sites in either of these glycoprotein hormones based on the amino acid differences between the two β -subunits. This approach also applies equally to non-humans as well.

15

The present invention also provides a modified human glycoprotein hormone having increased activity over a wild-type human glycoprotein, wherein the modified human hormone comprises a basic amino acid substituted at a position corresponding to the same amino acid position in a non-human glycoprotein hormone having an increased activity over the wild-type human glycoprotein.

20

The non-human glycoprotein hormone having an increased activity over the wild-type human glycoprotein can be any non-human species. For example, the non-human species can be bovine. See, for example, *Benua, R.S., et al.* "An 18 year study of the use of beef thyrotropin to increase I-131 uptake in metastatic thyroid cancer." J. Nucl. Med. 5:796-801 (1964) and *Hershman, J.M., et al.* Serum thyrotropin (TSH) levels after thyroid ablation compared with TSH levels after exogenous bovine TSH: implications for I-131 treatment of thyroid carcinoma." J. Clin. Endocrinol. Metab. 34:814-818 (1972). Alternatively, the non-human species can be equine, porcine, ovine, and the like. In the Example contained herein, the sequence of the 10-21 amino acid region of 27 species is set forth.

30

The present invention also provides a modified glycoprotein hormone having increased activity over a wild-type glycoprotein hormone from the same species, wherein the modified glycoprotein hormone comprises a basic amino acid substituted at
5 a position corresponding to the same amino acid position in a glycoprotein hormone from another species having an increased activity over the wild-type glycoprotein hormone. Therefore the glycoprotein being modified to increase its activity can be from a non-human species. For example, one can compare porcine glycoprotein hormones to bovine glycoprotein hormones, design porcine glycoprotein hormones with amino acid
10 substitutions at positions where the porcine and the bovine sequences are different, construct porcine glycoprotein hormones with the selected changes, and administer the modified porcine glycoprotein hormone to porcine animals. Alternatively, the glycoprotein hormone being modified can be bovine.

15 The present invention also provides a modified glycoprotein hormone having increased activity over the wild-type glycoprotein hormone from the same species, wherein the modified glycoprotein hormone comprises a basic amino acid substituted at a position corresponding to the same amino acid position in a different glycoprotein hormone from the same species having an increased activity over the wild-type
20 glycoprotein hormone. For example, the β -subunits of human thyroid-stimulating hormone and human chorionic gonadotropin can be compared and amino acid substitutions to either of these β -subunits can be made based on any sequence divergence. Naturally, only those changes which generally increase or decrease the activity of the modified glycoprotein hormone are contemplated since the hormone
25 receptor specificity will still need to be retained. An example of such a β -subunit modification is set forth in the Examples contained herein, where basic amino acids were substituted at positions 58 and 63 of the human thyroid stimulating hormone based on sequence comparison between the human thyroid stimulating hormone and the human chorionic gonadotropin hormone.

Modification refers to the substitution of a non-basic amino acid at any particular position or positions of the wild-type glycoprotein with a basic amino acid. In a presently preferred embodiment of the present invention, these modifications comprise the substitution of lysine for a non-basic amino acid.

5

The effect of the modification or modifications to the wild-type glycoprotein hormone can be ascertained in any number of ways. For example, cyclic AMP (cAMP) production in cells transfected with the modified glycoprotein can be measured and compared to the cAMP production of similar cells transfected with the wild-type glycoprotein hormone. Alternatively, progesterone production in cells transfected with the modified glycoprotein can be measured and compared to the progesterone production of similar cells transfected with the wild-type glycoprotein hormone. Alternatively, the activity of a modified glycoprotein hormone can be determined from receptor binding assays, from thymidine uptake assays, or from T₄ secretion assays.

Specific examples of such assays for determining the activity of modified glycoprotein hormones is set forth in the Example section contained herein. One skilled in the art can readily determine any appropriate assay to employ to determine the activity of either a wild-type or a modified glycoprotein hormone.

In one embodiment of the present invention, the modified glycoprotein hormone has an activity which is increased over the activity of the wild type glycoprotein hormone by at least 3 fold. This increased activity can be assessed by any of the techniques mentioned above and described in the Example contained herein, or in any other appropriate assay as readily determined by one skilled in the art. The increased activity does not have to be consistent from assay to assay, or from cell line to cell line, as these of course, will vary. For example, and as set forth in the Example contained herein, the relative potency of the P16K mutation in the α -subunit of the human glycoprotein hormone compared to the activity of the wild type glycoprotein hormone in a cAMP assay was approximately 6.4 fold higher. In the progesterone release assay, however, the difference between the same mutant and the wild-type glycoprotein hormone was approximately 3.4 fold in potency and 1.6 fold in V_{max}. This specific

modification demonstrates at least a 3 fold increase in activity in at least one assay, and therefore represents a glycoprotein hormone with at least a 3 fold increase in activity.

To modify additional amino acid positions, glycoprotein hormone sequences
5 from human and non-humans can be aligned using standard computer software
programs such as DNASIS (Hitachi Software Engineering Co. Ltd.). The amino acid
residues that differ between the human and the non-human glycoprotein hormone can
then be substituted using one of the above-mentioned techniques, and the resultant
glycoprotein hormone assayed for its activity using one of the above-mentioned assays.

10

The subject being treated or administered a modified glycoprotein hormone can
be a human or any non-human mammal. For example, the modified glycoprotein
hormone superagonists may be used in the superovulation of bovine animals by
administering these glycoprotein hormones to those bovine animals.

15

The methods used in substituting a basic amino acid for a non-basic amino acid
at any particular position or positions can also be used to design glycoprotein hormone
antagonists. By making specific substitutions and monitoring the activity of these
modified glycoprotein hormones, one can determine which modifications yield
20 glycoprotein hormones with reduced activity. These glycoprotein hormone agonists can
be used in studies of the hormone receptor such as receptor turnover rates, receptor
affinity for the glycoprotein hormone, or even in therapeutic procedures such as
treatment of Grave's disease and in fertility control.

25 The present invention also provides a method of treating a condition associated
with a glycoprotein hormone activity in a subject comprising administering a therapeutic
amount of the glycoprotein hormone of the present invention to the subject. These
conditions include any condition associated with a glycoprotein hormone activity.
Examples of these conditions include, but are not limited to, ovulatory dysfunction, luteal
30 phase defect, unexplained infertility, male factor infertility, time-limited conception.

In another example, the glycoprotein hormone may be administered to diagnose and treat a thyroid carcinoma. For example, the administration of bovine TSH to a human subject can be used to stimulate the uptake of ^{131}I in thyroid tissue to treat thyroid carcinoma. (*Meier, C.A., et al.*, "Diagnostic use of Recombinant Human Thyrotropin in Patients with Thyroid Carcinoma (Phase I/II Study)." *J. Clin. Endocrinol. Metabol.* 78:22 (1994)).

A skilled practitioner in the art can readily determine the effective amount of the glycoprotein hormone to administer and will depend on factors such as weight, size, the severity of the specific condition, and the type of subject itself. The therapeutically effective amount can readily be determined by routine optimization procedures. The present invention provides glycoprotein hormones with increased activity relative to the wild-type glycoprotein hormone. These modified glycoprotein hormones will allow a skilled practitioner to administer a lower dose of a modified glycoprotein hormone relative to the wild-type glycoprotein hormones to achieve a similar therapeutic effect, or alternatively, administer a dose of the modified glycoprotein hormone similar to the dose of the wild-type glycoprotein hormone to achieve an increased therapeutic effect.

Depending on whether the glycoprotein hormone is administered orally, parenterally, or otherwise, the administration of the prostaglandin can be in the form of solid, semi-solid, or liquid dosage forms, such as, for example, tablets, pills, capsules, powders, liquids, creams, and suspensions, or the like, preferably in unit dosage form suitable for delivery of a precise dosage. The glycoprotein hormone may include an effective amount of the selected glycoprotein hormone in combination with a pharmaceutically acceptable carrier and, in addition, may include other medicinal agents, pharmaceutical agents, carriers, adjuvants, diluents, etc. By "pharmaceutically acceptable" is meant a material that is not biologically or otherwise undesirable, i.e., the material may be administered to an individual along with the selected glycoprotein hormone without causing unacceptable biological effects or interacting in an unacceptable manner with the glycoprotein hormone. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see

Remington's Pharmaceutical Sciences, latest edition (Mack Publishing Co., Easton, PA.).

In another aspect, the present invention provides a method of assisting reproduction in a subject comprising administering an assisting amount of the glycoprotein hormone of the present invention. For example, in a subject with isolated gonadotropin deficiency (IGD), administration of modified follicle stimulating hormone (follitropin) and luteinizing hormone (lutropin) may be administered to the subject to restore normal gonadal function. It is widely known in the art that glycoprotein hormones such as FSH and LH are integral in female reproductive physiology, and these glycoprotein hormones may be administered to a subject to overcome a number of reproductive disorders and thereby assist reproduction.

Genetic therapy is another approach for treating hormone disorders with the modified glycoprotein hormones of the present invention. In this approach, a gene encoding the modified glycoprotein hormone can be introduced into a cell, such as a germ line cell or a somatic cell, so that the gene is expressed in the cell and subsequent generations of those cells are capable of expressing the introduced gene. For example, any particular gonadotropin hormone can be inserted into an ovarian cell, or its precursor, to enhance ovulation. Alternatively, introducing thyroid cells carrying a gene encoding a superagonist of the thyroid stimulating hormone into an individual with thyroid carcinoma can obviate the need for continual administration of TSH for stimulating radioiodine uptake in the thyroid carcinoma. Suitable vectors to deliver the coding sequence are well known in the art. For example, the vector could be viral, such as adenoviral, adenoassociated virus, retrovirus, or non-viral, such as cationic liposomes.

The modified glycoprotein hormones as provided by the present invention can also be used for targeting delivery of therapeutic agents to thyroid tissues or gonadal tissue, or in the treatment of certain neoplasms.

In yet another aspect, the invention provides a method of constructing superactive nonchimeric analogs of human hormones comprising comparing the amino acid sequence of a more active homolog from another species to the human hormone, substituting amino acids in the human hormone with the corresponding amino acids from the other species, determining the activity of the substituted human hormone, and selecting superactive analogs from the substituted human hormones. Superactive analogs of human hormones includes any analog whose activity is increased over the corresponding activity of the wild-type hormone. For example, the modification of the human thyroid stimulating hormone at position 11 in the α -subunit from threonine to lysine (T11K) results in a relative increase in the cAMP production in JP09 cells cultured *in vitro*. (See Table II as set forth in the Example contained herein). This modification of the human thyroid stimulating hormone therefore results in a superactive analog of the wild-type human thyroid stimulating hormone. The specific amino acid or amino acids to substitute to create the modification can be determined, as discussed above, by: determining the activity of the homolog from another species and comparing that activity to the human hormone; then comparing the aligned sequences to determine the amino acid sequence differences; then substituting the appropriate amino acid in the hormone from another species for the amino acid at the corresponding position in the human hormone; then determining the activity of the modified human hormone by one of the above-mentioned techniques; and then comparing the activity of the modified human hormones to the wild-type human hormone, thereby selecting the superactive analogs from the substituted human hormones.

All combinations of amino acid substitutions may be utilized to obtain a glycoprotein superagonist. For example, neutral amino acids can be substituted for basic or acidic amino acids. Alternatively, basic amino acids can be substituted for acidic or neutral amino acids, or acidic amino acids may be substituted for neutral or basic amino acids. One skilled in the art will recognize, as discussed above, that substitution of one amino acid for another can be at either the nucleic acid level in the nucleotide sequence that encodes the glycoprotein hormone or part of the glycoprotein hormone, or at the polypeptide level. Any human hormone can be modified by this method and its

superactive analogs selected. In particular, the human hormone can be a glycoprotein hormone.

EXAMPLES

5 The sequence between Cys10 and Pro21 of the human α -subunit was selected as the primary target for mutagenesis (Fig. 1). hCG-based homology modeling suggested that this region of the α -subunit is distant from the β -subunit in all glycoprotein hormones, contains several surface-exposed residues and includes a single turn of a 3_{10} -helix between Pro16 and Ser19¹. The human α -subunit differs from bovine in position
10 11, 13, 16, 17 and 20 (Fig. 1a) and four of these changes are nonconservative (Thr11→Lys, Gln13→Lys, Pro16→Lys and Gln20→Lys). We used PCR amplification to determine the sequence of the 11-20 region in the α -subunit of several primates including higher apes (common chimpanzee - *Pan troglodytes*, orangutan - *Pongo pygmaeus*), lesser apes (gibbon - *Hylobates* sp.), Old World monkey (baboon - *Papio anubis*) and compare them with previously known mammalian sequences including
15 rhesus macaque (*Macaca mulatta*; Old World monkey), common marmoset (*Callithrix jacchus*; New World monkey) and human (Fig. 1a). Simultaneous comparison of the sequences between different species suggested that basic residues in this region were replaced relatively late in primate evolution. The Rhesus monkey α -subunit gene codes
20 for Lys residues at positions 11, 16 and 20 and an Arg residue at position 13², the baboon sequence codes for Gln at position 16, whereas gibbon sequence contains only one weakly basic imidazolium group of His at position 13 (Fig. 1a). Apparently a cluster of positively charged amino acids in this region was maintained and modified during vertebrate evolution, but is not present in the higher apes and human sequence.
25 The gradual elimination of positively charged residues in the 11-20 region of α -subunit coincide with the evolutionary divergence of the hominoids (human and apes) from the Old World monkeys. Our hypothesis that this region may modulate binding to the receptor was further supported by: 1) the highest reactivity of Tyr21 in bTSH toward iodination³, 2) mapping of antigenic determinants in hCG⁴, 3) the role of amino groups
30 of Lys in the ovine and human α -subunit for effective hormone-receptor interaction as

studied by acylation⁵, labeling with acetic anhydride⁶ and pegylation of individual subunits.

Consequently, positively charged Lys residues were inserted into the
5 Cys10-Pro21 region of the human α -subunit (Fig. 1b). Two other regions were also mutagenized (Table I). A single nonconservative Leu69→Arg mutation in the TSH β -subunit was made based on a similar sequence comparison.

Effects of mutations

10 Cotransfection of wild-type (WT) or mutant human α and hTSH β ⁷ or hCG β cDNAs in various combination into CHO-K1 cells resulted in the expression of 14 hTSH and 11 hCG heterodimers (Table I). In contrast to many other mutagenesis studies^{8,9} the expression of mutants was generally comparable to the WT. The following hTSH α -mutants were expressed at levels higher than WT-hTSH: T11K, Q13K, P16K,
15 Q20K, Q50P and Q13K+P16K+Q20K. Thus, this set of evolutionary justified mutations did not impair, in a major way, synthesis of the hTSH or hCG molecule, but may facilitate in certain cases hormone production.

Various bioassays were used to compare the relative potency and efficacy of hTSH
20 and hCG mutants. The ability of WT and mutant hTSH to stimulate cAMP production was tested in CHO-JP09 cells with stably transfected human TSH receptor. This assay revealed the following order of potencies in single α -subunit mutants: P16K (6-fold lower EC₅₀ than WT) \geq Q20K > Q13K > T11K > WT-hTSH \approx Q50P \approx R67K (Table II). Receptor binding activity of WT and mutants hTSH was assessed in a competitive
25 binding assay to porcine thyroid membranes. Consistent with the cAMP stimulation, the following order of potencies was observed: P16K (5-fold greater affinity than WT) > Q20K \geq Q13K > T11K > WT-hTSH \approx Q50P \approx R67K (Table II). Thus, the increase in potency of single mutants observed in JP09 cells was directly correlated with the increase of affinity to the TSH receptor. Most notably, each mutation to a Lys residue
30 in the 11-20 region caused a substantial increase in activity, but changes outside this critical region had no (R67K, Q50P) effect on receptor binding affinity and bioactivity

(Table II). Alanine mutagenesis of amino acids 13, 16 and 20 in hTSH did not significantly alter hormone activity, indicating that only selective reconstitution of basic amino acids present in homologous hormones of other species resulted in the functional changes. Moreover, the exchange of α Ser43 to Arg and the replacements of α His90 and α Lys91 showed that these residues were less important for hTSH than for hCG bioactivity, emphasizing hormone- and site-specific roles of basic residues⁹.

Superagonists with combined mutations

To further study the effect of Lys residues which were individually responsible for highest increases in potency, mutants containing multiple replacements were produced. The most active mutants are presented in Fig. 2 and 3. The double Pro16-Lys + Gln20-Lys and the triple Pro16-Lys + Gln20-Lys + Gln13-Lys mutants showed, respectively, 12 and 24-fold higher activity than WT-hTSH, with a further increase in potency up to 35-fold after Leu69-Arg replacement in the TSH β -subunit (Fig. 2a). Additional optimization included substitution Glu14-Lys (Lys in this position present in the tuna sequence) resulted in further increase in bioactivity up to 95-fold; these most potent multiple mutants elevated efficacy (maximal response) at least 1.5-fold (Fig. 2b). These increases were verified by testing the ability of hTSH mutants to bind to porcine as well as human TSH receptor (Table II, Fig. 2c and Fig. 2d), to induce growth in FRTL-5 cells (Fig. 2e), as well as T₃ production in cultured human thyroid follicles. In particular, Pro16-Lys + Gln20-Lys + Gln13-Lys / WT-hTSH β and Pro16-Lys + Gln20-Lys + Gln13-Lys / Leu69-Arg mutants required, respectively, 18- and 27-fold lower concentration to attain half-maximal stimulation of ³H-thymidine incorporation in FRTL-5 cells than the WT-hTSH (Fig. 2e). The synergistic effect of multiple mutations on TSH bioactivity was not limited to a local cooperation of Lys residues in the 13-20 region of the α -subunit with receptor, but also involved the contribution of Arg69 in the opposite loop of β -subunit (Table II).

Table I.**Relative expression of wild-type (WT) and mutant hormones in CHO-K1 cells**

		hTSH	hCG
5	WT	100 ± 7	100 ± 4
	T11K	267 ± 22	82 ± 2
	Q13K	188 ± 9	106 ± 7
	P16K	206 ± 25	72 ± 6
10	Q20K	149 ± 18	117 ± 8
	P16K+Q20K	86 ± 9	62 ± 6
	Q13K+P16K+Q20K	134 ± 6	76 ± 12
	Q13K+P16K+Q20K+E14K	76 ± 12	52 ± 8
	P16K+F17T	23 ± 10	93 ± 4
15	Q50P	174 ± 15	83 ± 3
	R67K	171 ± 14	88 ± 6
	β-L69R	74 ± 5	n.a.
	Q13K+P16K+Q20K+β-L69R	86 ± 6	n.a.
	Q13K+P16K+Q20K+E14K+β-L69R	25 ± 6	n.a.
20			

n.a. = not applicable. Secretion levels are given as mean ± SEM relative to the WT, which was defined as 100% of WT-hTSH or WT-hCG respectively. The mean was calculated from at least four independent transfections, performed in at least five dishes for each mutant.

- 25 These findings were further confirmed in the animal model. A single injection of Pro16→Lys, Gln20→Lys and Gln13→Lys hTSH mutants in mice increased serum T₄ significantly higher than the WT-hTSH. Moreover, Pro16→Lys + Gln20→Lys + Gln13→Lys / WT-hTSHβ and Pro16→Lys + Gln20→Lys + Gln13→Lys / Leu69→Arg mutants also generated higher T₄ levels as compared to WT-hTSH (Fig. 2f). hTSH
- 30 serum levels 6 h after i.p. injection in mice were similar and the hTSH analogs did not show compared to the WT great differences in the metabolic clearance rate.

Table II. The effects of site-specific mutagenesis of human glycoprotein hormones

	hTSH		hTCG	
	cAMP stimulation in JP09 cells EC ₅₀ (ng/ml)	Relative potency (WT=1)	Inhibition of ¹²⁵ I-bTSH binding (EC ₂₅ , ng/ml)	Progesterone synthesis in MA10 cells (EC ₅₀ , ng/ml; Max, %)
WT	6.70 ± 0.69	1.0	81.3 ± 13.8	6.90 ± 1.04
T11K	4.47 ± 0.79	1.5	68.3 ± 4.4	2.79 ± 0.25
Q13K	1.89 ± 0.41	3.5	22.5 ± 2.6	2.46 ± 0.28
P16K	1.05 ± 0.26	6.4	18.3 ± 3.6	2.05 ± 0.17
Q20K	1.16 ± 0.22	5.8	21.3 ± 3.8	2.98 ± 0.27
P16K+Q20K	0.57 ± 0.10	11.8	6.4 ± 2.4	1.70 ± 0.13
Q13K+P16K+Q20K	0.28 ± 0.07	23.9	2.3 ± 0.3	1.58 ± 0.09
Q13K+P16K+Q20K +E14K	0.17 ± 0.04	39.4	2.1 ± 0.4	1.65 ± 0.06
P16K+FTT	3.52 ± 0.50	1.9	n.d.	n.d.
Q50P	5.54 ± 0.70	1.2	77.5 ± 12.4	3.90 ± 0.85
R67K	7.36 ± 0.33	0.9	62.5 ± 15.5	4.60 ± 0.63
TSHβ-L69R	2.75 ± 0.49	2.4	n.d.	n.a.
TSHβ-L69R+ +Q13K+P16K+Q20K	0.19 ± 0.06	35.3	1.8 ± 0.3	n.a.
TSHβ-L69R+ +Q13K+P16K+Q20K	0.07 ± 0.02	95.7	1.3 ± 0.4	n.a.
bTSH	0.71 ± 0.14	9.4	7.9 ± 2.5	n.a.

n.s. - no stimulation; n.a. - not applicable; n.d. - not determined. A curve-fitting program, Mac Allfit (NIH, Bethesda, MD) was used to fit the dose-response data and calculate EC₅₀ and Vmax values. Values are the mean ± SEM. Pro 16/Lys + Phe 17/Thr mutant with glycosylation consensus sequence (Asn-Lys-Thr) was created to study the effect of neoglycosylation on hormone activity.

A sequence comparison of the hCG and hTSH β -subunits showed a region (residues 58-69 in TSH β) which contains a cluster of basic residues in hCG, but not in hTSH. We used site-directed mutagenesis to introduce single and multiple basic residues into hTSH, based on their location in hCG, generating the additional
5 hTSH β -subunit mutants: I58R, E63R, I58R+E63R, I58R+E63R+L69R. The mutant hTSH β -subunits were coexpressed with the human α -subunit and the intrinsic activity of the recombinant hTSH analogs studied at the rat TSH receptor (FRTL-5 cells) and human TSH receptor (CHO-hTSHr cells). In both systems, single substitutions (I58R, E63R) increased potency of hTSH 2-fold to 4-fold, and
10 led to a slight increase of efficacy (Fig. 2g). The combination of the two substitutions (I58R+E63R) resulted in the potency which was 15-fold higher than that of wild type hTSH and an 1.5-fold increase of efficacy (Fig. 2g). Potency and efficacy of the combination mutant I58R+E63R+L69R, in which three basic residues were introduced, was elevated 50-fold and 1.7-fold, respectively (Fig. 2g).
15 These increases of intrinsic activity were accompanied by concomitant increases in receptor binding affinity, judged by a receptor-binding assay using CHO-JP09 cells (Fig. 2h). Similarly, when mice were injected with the I58R+E63R+L69R mutant, their T_4 stimulation was significantly higher than in either mock or control treated mice. (Fig. 2i).

20

The bioactivity of hCG mutants was tested using progesterone stimulation in MA-10 cells and cAMP stimulation in COS-7 cells transfected with human LH/hCG receptor. hCG Lys mutants showed both higher potency (lower EC_{50} values) as well as higher efficacy (V_{max}) than the WT-hCG (Table II, Fig. 3a). The
25 effect of single and multiple mutations was relatively analogous to that observed for the hTSH mutants. The α Pro16-Lys / WT-hCG β mutant was 4-fold more active than WT-hCG in the stimulation of progesterone production and receptor binding activity in MA-10 cells, with further increases in both potency and efficacy for the Pro16-Lys + Gln20-Lys and Pro16-Lys + Gln20-Lys hCG + Gln13-Lys
30 mutants (Fig. 3a and 3b). Similar increases of intrinsic activity were also found

when studied at the human LH/hCG receptor (COS-7-hLH/hCG-R cells) (Fig. 3c and 3d).

Our data suggest that only a few amino acid replacements are sufficient to
5 increase glycoprotein hormone bioactivity, even to a level higher than that of the
model hormone (such as bTSH). Interestingly, only a few of the 40 differing
residues between bovine and human TSH appear responsible for the higher
biological activity of bTSH. The majority of the other replacements are
conservative, and as illustrated by the R67K mutation in the α -subunit, seem to
10 have no functional significance. In contrast, we show that surface-located Lys
residues clustered in the L1 loop and β 1-strand of the α -subunit are crucial for the
high bioactivity of bTSH. Accordingly, recombinant hTSH with only two mutated
amino acids (P16K+Q20K) attains an intrinsic activity comparable to bTSH (Table
II). Moreover, triple, quadruplicate and quintuple hTSH mutants show even higher
15 potency than bTSH. These data suggest that the difference in activity between
bTSH and hTSH is a result of several amino acid changes, including replacements
increasing activity, but also others which may reduce biopotency of bTSH at the
hTSH receptor.

20 Although, we cannot exclude a possibility that several receptor species would
be made from a single transfected cDNA (by alternative splicing from cryptic sites
or by posttranslational modifications), the fact that similar differences in activity
were observed in different cell systems argues strongly against the importance of
different receptor species in the increase in potency, efficacy and affinity of these
25 analogs. Furthermore, there is compelling evidence that naturally occurring
hormone isoforms with various carbohydrate residues exert their effect at the post-
receptor level with no or minimal effect on receptor binding affinity¹⁰. Since the
wild type hormones and their analogs were characterized in multiple experimental
systems, it is highly probable that phenomenon of increased bioactivity described
30 here is a rule rather than exception related to particular cell-dependent variant of
the receptor.

Perspectives of rational design of glycoprotein hormone analogs

Previous site-directed mutagenesis studies of glycoprotein hormones focused primarily on the highly conserved regions and residues, using such strategies as alanine scanning mutagenesis¹¹ or multiple replacement approaches⁹. Several important studies were based on the creation of chimeric subunits using cassette mutagenesis and/or restriction fragment exchange^{12,13,14}. Our strategy based on replacement of nonconserved residues to those present in other species has been successful and permitted the generation of other glycoprotein hormone analogs, including hFSH mutants with increased bioactivity. The parallel improvement of bioactivity of hTSH, hCG and hFSH by introduction of basic residues in the 11-20 region of human α -subunit may be related to the fact that this region is distant from the β -subunit in the crystal structure based model of hCG and in our homology model of hTSH. The virtual identity of this area in both models as well as the observation that the antibodies binding to 11-26 region are not greatly influenced by subunit combination¹⁵ suggest that this domain may function similarly in all the glycoprotein hormones. Once the α -subunit was successfully engineered to create more potent agonists of hTSH, hCG or hFSH, the same paradigm was used to modify their respective β -subunits to generate the ultimate superagonists of each glycoprotein hormone. For example, an additional replacement of a nonpolar Leu69 to Arg in the TSH β -subunit resulted in further increase of hTSH bioactivity. In addition, the plasma half-life of our analogs can be modified regarding to specific therapeutic needs.

Further design and refinement of glycoprotein hormone analogs will include detailed three-dimensional structure of the hormone-receptor complexes. Although the exact structure of glycoprotein hormone receptors has not been solved, several models of hormone-receptor interaction have been proposed^{15,16,17,18}. In accordance with the recent model of Jiang et al.¹⁷ the L1 loop of α -subunit may participate in the interaction with the transmembrane portion of the receptor. The cluster of positively charged residues in this loop may enhance

such an interaction and facilitate further rearrangements in the receptor leading to the activation of G proteins and signal transduction.

Methods and Materials.

5 Restriction enzymes, DNA markers and other molecular biological reagents were purchased from either Gibco BRL (Gaithersburg, MD) or from Boehringer-Mannheim (Indianapolis, IN). Cell culture media, fetal bovine serum and LipofectAMINE were purchased from Gibco BRL (Gaithersburg, MD). VentR DNA Polymerase was purchased from New England Biolabs (Beverly, MA). The
10 full length human α cDNA (840 bp) subcloned into BamHI/XhoI sites of the pcDNA I/Neo vector (Invitrogen Corp., San Diego, CA) and hCG- β gene were obtained from Dr. T.H. Ji (University of Wyoming, Laramie, WA). The hTSH- β minigene without the first intron, with the non-translated 1st exon and authentic translation initiation site was constructed in our laboratory. rhTSH-G standard was
15 from Genzyme Corp. (Framingham, MA). The CHO cells with stably expressed hTSH receptor (CHO-hTSHR clone JP09 and clone JP26) were provided by Dr. G. Vassart (University of Brussels, Brussels, Belgium). The human LH receptor cDNA was obtained from Dr. T. Minegishi (Gunma University, Gunma, Japan). FRTL-5 cells were kindly supplied by Dr. L.D. Kohn (NIDDK, NIH, Bethesda,
20 MD). MA-10 cells were generously supplied by Dr. M. Ascoli (University of Iowa, Iowa City, IA). ^{125}I cAMP and ^{125}I -hTSH were from Hazleton Biologicals (Vienna, VA). Blood samples of various primates were obtained from Yerkes Regional Primate Research Center (Emory University, Atlanta, GA) and Animal Resources (University of Oklahoma, Oklahoma City, OK).

25 Determination of primate α -subunit sequences

. The QIAamp^R Blood Kit (Qiagen Inc., Chatsworth, CA) was used for extraction of genomic DNA from whole blood samples of chimpanzee (*Pan troglodytes*), orangutan (*Pongo pygmaeus*), gibbon (*Hylobates* sp.) and baboon
30 (*Papio anubis*). Genomic DNA was used in the PCR; the synthetic oligonucleotide

primers used were

5'-CCTGATAGATTGCCCAGAATGC-3' (sense) (SEQ ID NO:1) and

5'-GTGATAATAACAAGTACTGCAGTG-3' (antisense) (SEQ ID NO:2)

and were synthesized according to the nucleotide sequence of the gene encoding
5 common α -subunit of human glycoprotein hormones¹⁹. PCR was performed using
800-1000 ng of genomic DNA template and 10 picomoles of each primer in 100 μ l
reaction volume that also contained 10 mM Tris-HCl, (pH 9.0 at 25°C), 50 mM
KCl, 2.5 mM MgCl₂, 200 μ M dNTPs and 2 U of Taq DNA Polymerase (Promega
Corp. Madison, WI). The reaction mix was covered with mineral oil, and each
10 sample was initially heated to 95°C for 10 min. The PCR program consisted of 32
cycles of denaturation at 95°C for 1 min 30 sec, annealing at 55°C for 1 min 30 sec
and extension at 72°C for 1 min, followed by a final extension period at 72°C for 7
min. The reactions were then directly electrophoresed on a 1% agarose gel in the
presence of ethidium bromide. The amplified PCR product (~700 bp), spanning the
15 nucleotide sequence of exon 3, intron 3 and exon 4, was purified using QIAquick
PCR Purification Kit (QIAGEN Inc., Chatsworth, CA) and subcloned into
pCRTMII using Original TA Cloning Kit (Invitrogen Corp., San Diego, CA). The
sequence of the fragment was obtained after subcloning or direct dideoxy
sequencing using a Sequenase kit (U.S. Biochemical Corp., Cleveland, OH).

20 Homology modeling.

Modeling relies on the strong sequence homology between hCG and hTSH.
The sequences were aligned to bring the cysteine-knot residues into
correspondence and the percentage of identical as well as highly conservative
25 replacements were calculated as described¹. There was 58% sequence identity
between hCG and hTSH molecules; 31% of the two β -subunit sequences were
identical and additional 17% included highly conservative changes in β -subunit. A
molecular model of hTSH was built on a template of hCG model derived from
crystallographic coordinates obtained from the Brookhaven Data Bank²⁰. All
30 coordinate manipulations and energy calculations were done using CHARMM

release 21.2 for the Convex and further modified using the molecular graphic package QUANTA (Version 3.3, Molecular Simulations Inc., University of York, York, United Kingdom).

Site-directed mutagenesis.

5

- Mutagenesis of the human α -cDNA and the hTSH β minigene was accomplished by the PCR-based megaprimer method²¹. Amplification was optimized using Vent^R DNA Polymerase (New England Biolabs, Beverly, MA). After digestion with BamHI and XhoI PCR product was ligated into pcDNA I/Neo (Invitrogen Corp., San Diego, CA) with the BamHI/XhoI fragment excised. MC1061/p3 E. coli cells were transformed using Ultracomp E. coli Transformation Kit (Invitrogen Corp.). The QIAprep 8 Plasmid Kit (QIAGEN Inc., Chatsworth, CA) was used for multiple plasmid DNA preparations. QIAGEN Mega and Maxi Purification Protocols were used to purify larger quantities of plasmid DNA.
- 15 Multiple mutants were created with the same method using plasmids containing α -cDNA with a single mutation as a template for further mutagenesis. Mutations were confirmed by double stranded sequencing using Sanger's dideoxynucleotide chain termination procedure.

Expression of recombinant

20 **hormones.**

- CHO-K1 Cells (ATCC, Rockville, MD) were maintained in Ham's F-12 medium with glutamine and 10% FBS, penicillin (50 units/ml) and streptomycin (50 μ g/ml). Plates of cells (100 mm culture dishes) were cotransfected with wild type or mutant α -cDNA in the pcDNA I/NEO and hTSH β minigene inserted into the p(LB)CMV vector⁷, or pcDNAI/Neo containing hCG β -cDNA⁸ using a LipofectAMINE (Gibco BRL, Gaithersburg, MD). After 24 h, the transfected cells were transferred to CHO-serum free medium (CHO-SFM-II, Gibco BRL,). The culture media including control medium from mock transfections using the expression plasmids without gene inserts were harvested 72 h after transfection,
- 30

concentrated and centrifuged; the aliquots were stored at -20°C and thawed only once before each assay. WT and mutant hTSH were measured and verified using four different immunoassays as described⁹. Concentrations of WT and mutant hCG were measured using chemiluminescence assay (hCG Kit, Nichols Institute, San Juan Capistrano, CA) and immunoradiometric assay (hCG IRMA, ICN, Costa Mesa, CA).

cAMP stimulation in JP09 cells expressing human TSH receptor.

CHO cells stably transfected with hTSH receptor cDNA (JP09) were grown and incubated with serial dilutions of WT and mutant hTSH as described⁹. cAMP released into the medium was measured by radioimmunoassay²². The equivalent amounts of total media protein were used as the mock control and the hTSH containing samples from transfected cells.

cAMP stimulation in COS-7 cells expressing human LH receptor.

COS-7 cells transiently transfected with hLH receptor cDNA were grown and incubated with serial dilutions of WT and mutant hCG essentially as described²³. cAMP released into the medium was measured by radioimmunoassay²². The equivalent amounts of total media protein were used as the mock control and the hCG containing samples from transfected cells.

Progesterone production stimulation in MA-10 cells.

Transformed murine Leydig cells (MA-10) grown in 96-well culture plates were incubated with WT and mutant hCG for 6 hours in the assay medium as described²⁴. The amount of progesterone released into the medium was determined by radioimmunoassay (CT Progesterone Kit, ICN Biomedicals, Inc., Costa Mesa, CA).

Receptor binding assays.

The receptor-binding activities of hTSH analogs were assayed by their ability to displace ^{125}I -bTSH from a solubilized porcine thyroid membranes²²⁴. The binding activities of selected analogs to human TSH receptor was tested using JP09 cells. The binding activities of hCG analogs to MA-cells and to COS-7 cells transiently transfected with human LH receptor were determined using ^{125}I -hCG and assay medium as described previously²⁴.

Thymidine uptake stimulation in FRTL-5 cells.

10

Growth of the rat thyroid cells (FRTL-5) was monitored as previously described²².

Stimulation of T_4 secretion in mice.

15

The *in vivo* bioactivity of the WT and mutant TSH was determined using a modified McKenzie bioassay^{22,25}. WT and mutant TSH were injected i.p. into male albino Swiss Crl:CF-1 mice with previously suppressed endogenous TSH by administration of 3 $\mu\text{g}/\text{ml}$ T_3 in drinking water for 6 days. Blood samples were collected 6 h later from orbital sinus and the serum T_4 and TSH levels were measured by respective chemiluminescence assays (Nichols Institute).

Throughout this application various publications are referenced. Certain publications are referenced by numbers within parentheses. Full citations for the number-referenced publications are listed below. The disclosures of all of these publications and those references cited within those publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

It will be apparent to those skilled in the art that various modifications and variations can be made in the present invention without departing from the scope or spirit of the invention. Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.

References

1. Wu, H., Lustbader, J.W., Liu, Y., Canfield, R.E. and Hendrickson W.A. Structure of human chorionic gonadotropin at 2.6 Å resolution from MAD analysis of the selenomethionyl protein. *Structure* 2:545-558 (1994).
2. Golos, T.G., Durning, M. and Fisher, J.M. Molecular cloning of the Rhesus glycoprotein hormone α -subunit gene. *DNA Cell. Biol.* 10:367-380 (1991).
3. Stanton, P.G. and Hearn, M.T.W. The iodination sites of bovine thyrotropin. *J. Biol. Chem.* 262:1623-1632 (1987).
4. Dirnhofer S. et al. Free α -subunit of human chorionic gonadotrophin: molecular basis of immunologically and biologically active domains. *J. Endocrinol.* 140:145-154 (1994).
5. Liu, W.K., Yang, K.P. and Ward, D.N. The role of the amino group in the subunit association and receptor site interaction for ovine luteinizing hormone as studied by acylation. *J. Biol. Chem.* 249:5544-5550 (1974).
6. Yadav, S.P., Brew, K. and Puett, D. Holoprotein formation of human chorionic gonadotropin: differential trace labeling with acetic anhydride. *Mol. Endocrinol.* 8:1547-1558 (1994).
7. Joshi, L. et al. Recombinant thyrotropin containing a β -subunit chimera with the human chorionic gonadotropin- β carboxy terminus is biologically active, with a prolonged plasma half-life: role of carbohydrate in bioactivity and metabolic clearance. *Endocrinology* 136:3839-3848 (1995).
8. Ji, I., Zeng H. & Ji, T.H. J. Receptor activation of and signal generation by the lutropin/choriogonadotropin receptor. *Biol. Chem.* 268:22971-22974 (1993).

9. Grossmann, M. et al. Role of the carboxy-terminal residues of the α -subunit in the expression and bioactivity of human thyroid-stimulating hormone. *Mol. Endocrinol.* 9:948-958 (1995).
10. Szkudlinski, M.W., Thotakura, N.R. and Weintraub, B.D. Subunit-specific functions of N-linked oligosaccharides in human thyrotropin: Role of terminal residues of α -subunit and β -subunit oligosaccharides in metabolic clearance and bioactivity. *Proc. Natl. Acad. Sci. USA* 92:9062-9066 (1995).
11. Liu, C., Roth, K.E., Lindau-Shepard, B.A., Shaffer, J.B. and Dias, J.A. Site-directed alanine mutagenesis of Phe³³, Arg³⁵, and Arg⁴²-Ser⁴³-Lys⁴⁴ in the human gonadotropin α -subunit. *J. Biol. Chem.* 269:25289-25294 (1994).
12. Lunardi-Iskandar, Y. et al. Tumorigenesis and metastasis of neoplastic Kaposi's sarcoma cell line in immunodeficient mice blocked by a human pregnancy hormone. *Nature* 375:64-68 (1995).
13. Campbell, R.K., Dean-Emig, D.M. and Moyle, W.R. Conversion of human choriogonadotropin into a follitropin by protein engineering. *Proc. Natl. Acad. Sci.* 88:760-764 (1991).
14. Dias, J.A., Zhang, Y. and Liu, X. Receptor binding and functional properties of chimeric human follitropin prepared by an exchange between a small hydrophilic intercysteine loop of human follitropin and human lutropin. *J. Biol. Chem.* 269:25289-25294 (1994).
15. Moyle, W.R. et al. Model of human chorionic gonadotropin and lutropin receptor interaction that explains signal transduction of the glycoprotein hormones. *J. Biol. Chem.* 270:20020-20031 (1995).

16. Combarous, Y. Molecular basis of the specificity of binding of glycoprotein hormones to their receptors. *Endocrine Rev.* 13:670-691 (1992).
17. Jiang, X. et al. Structural predictions for the ligand-binding region of glycoprotein hormone receptors and the nature of hormone-receptor interactions. *Structure* 3:1341-1353 (1995).
18. Kajava, A.V., Vassart, G. and Wodak, S.J. Modeling of the three-dimensional structure of proteins with the typical leucine-rich repeats. *Structure* 3:867-877 (1995).
19. Fiddes, J.C. and Goodman, H.M. Isolation, cloning and sequence analysis of the cDNA for the α -subunit of human chorionic gonadotropin. *Nature* 281:351-356 (1979).
20. Laphorn, A.J. et al. Crystal structure of human chorionic gonadotropin. *Nature* 369:455-461 (1994).
21. Sarkar, G. and Sommer, S.S. The "megaprimer" method of site-directed mutagenesis. *BioTechniques* 8:404-407 (1990).
22. Szkudlinski, M.W. et al. Purification and characterization of recombinant human thyrotropin isoforms produced by Chinese hamster ovary cells: the role of sialylation and sulfation in thyrotropin bioactivity. *Endocrinology* 133:1490-1503 (1993).
23. Igarashi, S. et al. Functional expression of recombinant human luteinizing hormone/human choriogonadotropin receptor. *Biochem. Biophys. Res. Commun.* 201:248-256 (1994).

24. Ascoli, M. Characterization of several clonal lines of cultured Leydig tumor cells: gonadotropin receptors and steroidogenic responses. *Endocrinology* 108:88-95 (1981).
25. Moyle, W. R. et al. Co-evolution of ligand-receptor pairs. *Nature* 268:251-255 (1994).

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: THE UNITED STATES OF AMERICA AS REPRESENTED BY THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES IN CARE OF NATIONAL INSTITUTES OF HEALTH Office of Technology Transfer Suite 325, 6011 Executive Boulevard, Rockville, MD 20852

(ii) TITLE OF INVENTION: GLYCOPROTEIN HORMONE SUPERAGONISTS

(iii) NUMBER OF SEQUENCES: 37

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: NEEDLE & ROSENBERG, P.C.
(B) STREET: Suite 1200, 127 Peachtree Street
(C) CITY: Atlanta
(D) STATE: Georgia
(E) COUNTRY: USA
(F) ZIP: 30303

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Perryman, David G.
(B) REGISTRATION NUMBER: 33,438
(C) REFERENCE/DOCKET NUMBER: 14014.0185/P

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 404/688-0770

47

(B) TELEFAX: 404/688-9880

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CCTGATAGAT TGCCCAGAAT GC

22

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GTGATAATAA CAAGTACTGC AGTG

24

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant

48

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Cys	Thr	Leu	Gln	Glu	Asn	Pro	Phe	Phe	Ser	Gln	Pro
1				5					10		

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Cys	Thr	Leu	Gln	Glu	Asn	Pro	Phe	Phe	Ser	Gln	Pro
1				5					10		

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

49

Cys Thr Leu Gln Glu Asn Pro Phe Phe Ser Gln Pro
1 5 10

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Cys Gln Leu His Glu Asn Pro Phe Phe Ser Gln Pro
1 5 10

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Cys Lys Pro Arg Glu Asn Gln Phe Phe Ser Lys Pro
1 5 10

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Cys	Lys	Pro	Arg	Glu	Asn	Lys	Phe	Phe	Ser	Lys	Pro
1				5						10	

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Cys	Lys	Leu	Lys	Glu	Asn	Lys	Tyr	Phe	Ser	Arg	Leu
1				5						10	

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

51

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Cys	Lys	Leu	Lys	Glu	Asn	Lys	Tyr	Phe	Ser	Lys	Pro
1			5							10	

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Cys	Lys	Leu	Lys	Glu	Asn	Lys	Tyr	Phe	Ser	Lys	Pro
1			5							10	

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Cys	Lys	Leu	Arg	Glu	Asn	Lys	Tyr	Phe	Phe	Lys	Leu
1				5						10	

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Cys Lys Leu Lys Glu Asn Lys Tyr Phe Ser Lys Leu
1 5 10

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Cys Lys Leu Lys Glu Asn Lys Tyr Phe Ser Lys Leu
1 5 10

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Cys Lys Leu Lys Glu Asn Lys Tyr Phe Ser Lys Leu
1 5 10

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: not relevant

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Cys Lys Leu Lys Glu Asn Lys Tyr Phe Ser Lys Leu
1 5 10

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: not relevant

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Cys Lys Leu Lys Gln Asn Lys Tyr Phe Ser Lys Leu
1 5 10

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Cys	Lys	Leu	Gly	Glu	Asn	Arg	Phe	Phe	Ser	Lys	Pro
1				5						10	

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Cys	Lys	Leu	Gly	Glu	Asn	Arg	Phe	Phe	Ser	Lys	Pro
1				5						10	

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

55

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Cys	Lys	Leu	Gly	Glu	Asn	Arg	Phe	Phe	Ser	Lys	Pro
1				5						10	

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Cys	Arg	Leu	Lys	Glu	Asn	Leu	Arg	Phe	Ser	Asn	Met
1				5						10	

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Cys	Lys	Leu	Lys	Glu	Asn	Lys	Val	Phe	Ser	Asn	Pro
1				5						10	

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Cys Thr Leu Lys Pro Asn Thr Ile Phe Pro Asn Pro
1 5 10

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Cys Lys Leu Lys Glu Asn Asn Ile Phe Ser Lys Pro
1 5 10

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

57

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Cys	Thr	Leu	Lys	Lys	Asn	Asn	Val	Phe	Ser	Arg	Asp
1				5						10	

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Cys	Thr	Leu	Arg	Lys	Asn	Thr	Val	Phe	Ser	Arg	Asp
1				5						10	

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Cys	Thr	Leu	Arg	Lys	Asn	Ser	Val	Phe	Ser	Arg	Asp
1				5						10	

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Cys	Lys	Leu	Lys	Glu	Asn	Asn	Ile	Phe	Ser	Lys	Pro
1				5						10	

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Cys	Thr	Leu	Lys	Glu	Asn	Asn	Ile	Phe	Ser	Lys	Pro
1				5						10	

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Cys Arg Leu Lys Asp Asn Lys Phe Phe Ser Lys Pro
1 5 10

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Cys Arg Leu Gln Glu Asn Lys Ile Phe Ser Lys Pro
1 5 10

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Cys Thr Leu Lys Glu Asn Pro Phe Phe Ser Gln Pro
1 5 10

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Cys	Thr	Leu	Gln	Glu	Asn	Lys	Phe	Phe	Ser	Gln	Pro
1				5						10	

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Cys	Thr	Leu	Gln	Glu	Asn	Pro	Phe	Phe	Ser	Lys	Pro
1				5						10	

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

61

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

Cys	Thr	Leu	Gln	Glu	Asn	Lys	Phe	Phe	Ser	Lys	Pro
1				5						10	

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Cys	Thr	Leu	Lys	Glu	Asn	Lys	Phe	Phe	Ser	Lys	Pro
1				5						10	

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Cys	Thr	Leu	Lys	Lys	Asn	Lys	Phe	Phe	Ser	Lys	Pro
1				5						10	

What is claimed is:

1. A human glycoprotein hormone comprising at least three basic amino acids in the α -subunit at positions selected from the group consisting of positions 11, 13, 14, 16, 17, and 20.
2. The human glycoprotein hormone of claim 1, further comprising a fourth basic amino acid at a position selected from the group consisting of positions 11, 13, 14, 16, 17, and 20.
3. The human glycoprotein hormone of claim 2, wherein basic amino acids are at positions 11, 13, 16, and 20.
4. The human glycoprotein hormone of claim 2, wherein basic amino acids are at positions 11, 13, 17, and 20.
5. The human glycoprotein hormone of claim 2, wherein basic amino acids are at positions 13, 14, 16, and 20.
6. The human glycoprotein hormone of claim 2, wherein basic amino acids are at positions 13, 14, 17, and 20.
7. The human glycoprotein hormone of claim 1, further comprising a fifth basic amino acid at a position selected from the group consisting of positions 11, 13, 14, 16, 17, and 20.
8. The human glycoprotein hormone of claim 7, wherein basic amino acids are at positions 13, 14, 16, 17, and 20.
9. The human glycoprotein hormone of claim 7, wherein basic amino acids are at positions 11, 13, 14, 16 and 20.
10. The human glycoprotein hormone of claim 1, wherein basic amino acids are at positions 11, 13, 14, 16, 17, and 20.

11. The human glycoprotein hormone of claim 1, wherein basic amino acids are at positions 13, 16, and 20.
12. The glycoprotein hormone of any of claims 1 - 11, wherein the hormone is thyroid stimulating hormone.
13. The human thyroid stimulating hormone of claim 12, wherein the hormone further comprises a basic amino acid in at least one position selected from the group consisting of positions 58, 63, and 69 of the β -subunit.
14. The human thyroid stimulating hormone of claim 13, wherein the hormone comprises basic amino acids at positions 58, 63, and 69 of the β -subunit.
15. The human glycoprotein hormone of claim 13, wherein a basic amino acid is at position 58.
16. The human glycoprotein hormone of claim 13, wherein a basic amino acid is at position 63.
17. The human glycoprotein hormone of claim 13, wherein a basic amino acid is at position 69.
18. The glycoprotein hormone of any of claims 1 - 11, wherein the hormone is follicle-stimulating hormone.
19. The glycoprotein hormone of any of claims 1 - 11, wherein the hormone is luteinizing hormone.
20. The glycoprotein hormone of any of claims 1 - 11, wherein the hormone is chorionic gonadotropin.
21. The human glycoprotein hormone of any of claims 1 - 20, wherein the basic amino acids are lysine.

22. The human glycoprotein hormone any of claims 1 - 20, wherein the basic amino acids are selected from the group consisting of lysine and arginine.
23. A method of making a medicament for treating a condition associated with a glycoprotein hormone activity in a subject comprising administering a therapeutic amount of a suitable glycoprotein hormone of any of the above claims to the patient.
24. The method of claim 23, wherein the condition is ovulatory dysfunction.
25. The method of claim 23, wherein the condition is a luteal phase defect.
26. The method of claim 23, wherein the condition is unexplained infertility.
27. The method of claim 23, wherein the condition is male factor infertility.
28. The method of claim 23, wherein the condition is time-limited conception.
29. The method of claim 23, wherein the condition is thyroid carcinoma.
30. A method of assisting reproduction in a subject comprising administering an assisting amount of the glycoprotein hormone of any of claims 1 - 22.
31. A nucleic acid encoding the human glycoprotein of any of claims 1 - 22.
32. The nucleic acid of claim 31 in a vector suitable for expressing the nucleic acid.
33. The vector of claim 32 in a host suitable for expressing the nucleic acid.
34. The glycoprotein hormone of any of claims 1 - 22, wherein the human glycoprotein has less than five amino acid substitutions in the α -subunit in positions other than positions 11, 13, 14, 16, 17, and 20.

35. The glycoprotein hormone of any of claims 1 - 22, wherein the human glycoprotein has less than four amino acid substitutions in the α -subunit in positions other than positions 11, 13, 14, 16, 17, and 20.
36. The glycoprotein hormone of any of claims 1 - 22, wherein the human glycoprotein has less than three amino acid substitutions in the α -subunit in positions other than positions 11, 13, 14, 16, 17, and 20.
37. The glycoprotein hormone of any of claims 1 - 22, wherein the human glycoprotein has less than two amino acid substitutions in the α -subunit in positions other than positions 11, 13, 14, 16, 17, and 20.
38. The glycoprotein hormone of any of claims 1 - 22, wherein the human glycoprotein has the complete amino acid sequence homology with the human glycoprotein hormone in positions other than positions 11, 13, 14, 16, 17, and 20 of the α -subunit.
39. The human glycoprotein of any of claims 18, 19, and 20, wherein the hormone comprises a basic amino acid in at least one position selected from the group consisting of positions in the β -subunit of glycoprotein hormones, corresponding to positions 58, 63, and 69 of the β -subunit of the human thyroid stimulating hormone.
40. A human glycoprotein hormone comprising a basic amino acid in the α -subunit in at least one position selected from the group consisting of positions 11, 13, 14, 16, 17 and 20.
41. The human glycoprotein hormone of claim 40, wherein a basic amino acid is at position 11.
42. The human glycoprotein hormone of claim 40, wherein a basic amino acid is at position 13.

43. The human glycoprotein hormone of claim 40, wherein a basic amino acid is at position 14.
44. The human glycoprotein hormone of claim 40, wherein a basic amino acid is at position 16.
45. The human glycoprotein hormone of claim 40, wherein a basic amino acid is at position 17.
46. The human glycoprotein hormone of claim 40, wherein a basic amino acid is at position 20.
47. The human glycoprotein hormone of any of claims 41 - 46, wherein the basic amino acid is lysine.
48. The human glycoprotein hormone of any of claims 41 - 46, wherein the basic amino acid is selected from the group consisting of lysine and arginine.
49. The human glycoprotein hormone of claim 40, comprising a basic amino acid in at least two positions selected from the group consisting of positions 11, 13, 14, 16, 17, and 20.
50. The human glycoprotein hormone of claim 49, wherein basic amino acids are at positions 16 and 20.
51. The human glycoprotein hormone of claim 49, wherein basic amino acids are at positions 16 and 13.
52. The human glycoprotein hormone of claim 49, wherein basic amino acids are at positions 20 and 13.
53. The human glycoprotein hormone of any of claims 49 - 52, wherein the basic amino acid is selected from the group consisting of lysine and arginine.

54. The glycoprotein hormone of any of claims 41 - 52, wherein the hormone is thyroid stimulating hormone.
55. The human thyroid stimulating hormone of claim 54, wherein the hormone further comprises a basic amino acid in at least one position selected from the group consisting of positions 58, 63, and 69 of the β -subunit.
56. The human thyroid stimulating hormone of claim 55, wherein the hormone comprises basic amino acids at positions 58, 63, and 69 of the β -subunit.
57. The human glycoprotein hormone of claim 55, wherein a basic amino acid is at position 58.
58. The human glycoprotein hormone of claim 55, wherein a basic amino acid is at position 63.
59. The human glycoprotein hormone of claim 55, wherein a basic amino acid is at position 69.
60. The glycoprotein hormone of any of claims 41 - 53, wherein the hormone is follicle-stimulating hormone.
61. The glycoprotein hormone of any of claims 41 - 53, wherein the hormone is luteinizing hormone.
62. The glycoprotein hormone of any of claims 41 - 53, wherein the hormone is chorionic gonadotropin.
63. A method of treating a condition associated with a glycoprotein hormone activity in a subject comprising administering a therapeutic amount of the suitable glycoprotein hormone of any of claims 1 - 22, 31 - 39, 40 - 62, and 71 - 84 to the patient.
64. The method of claim 63, wherein the condition is ovulatory dysfunction.

- 65. The method of claim 63, wherein the condition is a luteal phase defect.
- 66. The method of claim 63, wherein the condition is unexplained infertility.
- 67. The method of claim 63, wherein the condition is male factor infertility.
- 68. The method of claim 63, wherein the condition is time-limited conception.
- 69. The method of claim 63, wherein the condition is thyroid carcinoma.
- 70. A method of assisting reproduction in a subject comprising administering an assisting amount of the glycoprotein hormone of any of claims 40 - 62.
- 71. A nucleic acid encoding the human glycoprotein of any of claims 41 - 53.
- 72. The nucleic acid of claim 71 in a vector suitable for expressing the nucleic acid.
- 73. The vector of claim 72 in a host suitable for expressing the nucleic acid.
- 74. The glycoprotein hormone of any of claims 40 - 62, wherein the human glycoprotein has less than five amino acid substitutions in the α -subunit in positions other than positions 11, 13, 14, 16, 17, and 20.
- 75. The glycoprotein hormone of any of claims 40 - 62, wherein the human glycoprotein has less than four amino acid substitutions in the α -subunit in positions other than positions 11, 13, 14, 16, 17, and 20.
- 76. The glycoprotein hormone of any of claims 40 - 62, wherein the human glycoprotein has less than three amino acid substitutions in the α -subunit in positions other than positions 11, 13, 14, 16, 17, and 20.
- 77. The glycoprotein hormone of any of claims 40 - 62, wherein the human glycoprotein has less than two amino acid substitutions in the α -subunit in positions other than positions 11, 13, 14, 16, 17, and 20.

78. The glycoprotein hormone of any of claims 40 - 62, wherein the human glycoprotein has the complete amino acid sequence homology with the human glycoprotein hormone in positions other than positions 11, 13, 14, 16, 17, and 20 of the α -subunit.
79. The human glycoprotein of claim 40 - 62, wherein the hormone comprises a basic amino acid in at least one position selected from the group consisting of positions in the β -subunit of glycoprotein hormones other than the thyroid stimulating hormone, corresponding to positions 58, 63, and 69 of the β -subunit of the human thyroid stimulating hormone.
80. A modified human glycoprotein hormone having increased activity over a wild-type human glycoprotein hormone, wherein the modified human glycoprotein hormone comprises a basic amino acid substituted at a position corresponding to the same amino acid position in a non-human glycoprotein hormone having an increased activity over the wild-type human glycoprotein hormone.
81. A modified glycoprotein hormone having increased activity over a wild-type glycoprotein hormone from the same species, wherein the modified glycoprotein hormone comprises a basic amino acid substituted at a position corresponding to the same amino acid position in a glycoprotein hormone from another species having an increased activity over the wild-type glycoprotein hormone.
82. The glycoprotein hormone of claim 80, wherein the non-human glycoprotein hormone is from a bovine.
83. The glycoprotein hormone of claim 80, wherein the basic amino acid is lysine.
84. The glycoprotein hormone of claim 80, wherein the activity is increased by at least 3 fold.
85. A method of constructing superactive nonchimeric analogs of human hormones comprising:

- a) comparing the amino acid sequence of a more active homolog from another species to the human hormone;
 - b) substituting amino acids in the human hormone with the corresponding amino acids from the other species;
 - c) determining the activity of the substituted human hormone; and
 - d) selecting superactive analogs from the substituted human hormones.
86. A method of constructing antagonist nonchimeric analogs of human hormones comprising:
- a) comparing the amino acid sequence of a less active homolog from another species to the human hormone;
 - b) substituting amino acids in the human hormone with the corresponding amino acids from the other species;
 - c) determining the activity of the substituted human hormone; and
 - d) selecting antagonist analogs from the substituted human hormones.
87. The method of any of claims 85 or 86, wherein the hormone is a glycoprotein hormone.

1/14

Fig. 1a

	10	11	12	13	14	15	16	17	18	19	20	21
Human	Cys	Thr	Leu	Gln	Glu	Asn	Pro	Phe	Phe	Ser	Gln	Pro
Chimpanzee	---	---	---	---	---	---	---	---	---	---	---	---
Orangutan	---	---	---	---	---	---	---	---	---	---	---	---
Gibbon	---	Gln	---	His	---	---	---	---	---	---	---	---
Baboon	---	Lys	Pro	Arg	---	---	Gln	---	---	---	Lys	---
Rhesus	---	Lys	Pro	Arg	---	---	Lys	---	---	---	Lys	---
Marmoset	---	Lys	---	Lys	---	---	Lys	Tyr	---	---	Arg	Leu
Bovine	---	Lys	---	Lys	---	---	Lys	Tyr	---	---	Lys	---
Ovine	---	Lys	---	Lys	---	---	Lys	Tyr	---	---	Lys	---
Equine	---	Lys	---	Arg	---	---	Lys	Tyr	---	---	Lys	---
Porcine	---	Lys	---	Lys	---	---	Lys	Tyr	---	Phe	Lys	Leu
Rabbit	---	Lys	---	Lys	---	---	Lys	Tyr	---	---	Lys	Leu
Mouse	---	Lys	---	Lys	---	---	Lys	Tyr	---	---	Lys	Leu
Rat	---	Lys	---	Lys	---	---	Lys	Tyr	---	---	Lys	Leu
Whale	---	Lys	---	Lys	Gln	---	Lys	Tyr	---	---	Lys	Leu
Quail	---	Lys	---	Gly	---	---	Arg	---	---	---	Lys	---
Chicken	---	Lys	---	Gly	---	---	Arg	---	---	---	Lys	---
Turkey	---	Lys	---	Gly	---	---	Arg	---	---	---	Lys	---
Bullfrog	---	Arg	---	Lys	---	---	Leu	Arg	---	---	Lys	---
Salmon 1	---	Lys	---	Lys	---	---	Lys	Val	---	---	Asn	Met
Salmon 2	---	---	---	Lys	Pro	---	Thr	Ile	---	Pro	Asn	---
Catfish	---	Lys	---	Lys	---	---	Asn	Ile	---	---	Lys	---
Tuna	---	---	---	Lys	Lys	---	Asn	Val	---	---	Arg	Asp
Yellowfin	---	---	---	Arg	Lys	---	Thr	Val	---	---	Arg	Asp
Bass	---	---	---	Arg	Lys	---	Ser	Val	---	---	Arg	Asp
Carp 1	---	Lys	---	Lys	---	---	Asn	Ile	---	---	Lys	---
Carp 2	---	---	---	Lys	---	---	Asn	Ile	---	---	Lys	---
Pike eel	---	Arg	---	Lys	Asp	---	Lys	---	---	---	Lys	---
Eel	---	Arg	---	---	---	---	Lys	Ile	---	---	Lys	---

Fig. 1b

Q13K	---	---	---	Lys	---	---	---	---	---	---	---	---
P16K	---	---	---	---	---	---	Lys	---	---	---	---	---
Q20K	---	---	---	---	---	---	---	---	---	---	Lys	---
P16K+Q20K	---	---	---	---	---	---	Lys	---	---	---	Lys	---
P16K+Q20K+Q13K	---	---	---	Lys	---	---	Lys	---	---	---	Lys	---
P16K+Q20K+Q13K+E14K	---	---	---	Lys	Lys	---	Lys	---	---	---	Lys	---




Fig. 2a

3/14




Fig. 2b

4/14




Fig. 2c

5/14




Fig. 2d

6/14




Fig. 2e

7/14




Fig. 2f

8/14




Fig. 2g

9/14




Fig. 2h

10/14




Fig. 2i

11/14




Fig. 3a

12/14




Fig. 3b

13/14




Fig. 3c

14/14




Fig. 3d

INTERNATIONAL SEARCH REPORT

Int. l. Application No
PCT/US 96/06483

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/16 C07K14/59 A61K38/24

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07K C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 90 02812 A (APPLIED RESEARCH SYSTEMS ARS HOLDING) 22 March 1990 see the whole document ---	1-87
A	ANIMAL REPRODUCTION SCIENCE, vol. 33, 1993, pages 71-82, XP000571973 WILSON J M ET AL: "SUPEROVULATION OF CATTLE WITH A RECOMBINANT-DNA BOVINE FOLLICLE STIMULATING HORMONE" see the whole document --- -/--	1-87

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *I* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

20 January 1997

Date of mailing of the international search report

29. 01. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Moreau, J

INTERNATIONAL SEARCH REPORT

Int. Patent Application No.
PCT/US 96/06483

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BIOCHEMISTRY 31 (41). 1992. 10094-10098, XP002023077 LEINUNG M C ET AL: "SYNTHETIC ANALOGS OF THE CARBOXYL-TERMINUS OF BETA THYROTROPIN THE IMPORTANCE OF BASIC AMINO ACIDS IN RECEPTOR BINDING ACTIVITY." see the whole document ---	1-87
A	EP 0 404 458 A (BUNGE (AUSTRALIA) PROPRIETARY LIMITED) 27 December 1990 cited in the application see the whole document -----	1-87

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/06483

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Please see Further Information sheet enclosed.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/219

Remark : Although claims 30 and 63-70 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/06483

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9002812	22-03-90	AU-A- 4076589 EP-A- 0386203 GR-A- 89100541 JP-T- 3502525	02-04-90 12-09-90 31-10-90 13-06-91
EP-A-404458	27-12-90	CA-A- 2019239	19-12-90